

Title	Blasticidin S-resistance gene (bsr) : a novel selectable marker for mammalian cells
Sub Title	
Author	泉, 雅子(Izumi, Masako) 宮澤, 宏(Miyazawa, Hiroshi) 鎌倉, 高志(Kamakura, Takashi) 山口, 勇(Yamaguchi, Isamu) 遠藤, 豊成(Endo, Toyoshige) 花岡, 文雄(Hanaoka, Fumio)
Publisher	共立薬科大学
Publication year	1992
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.37 (1992. ) ,p.99- 99
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	<a href="https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000037-0099">https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000037-0099</a>

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

## Blasticidin S-Resistance Gene (*bsr*) : A Novel Selectable Marker for Mammalian Cells\*

Masako IZUMI,\*\*, Hiroshi MIYAZAWA,\*\* Takashi KAMAKURA\*\*, Isamu YAMAGUCHI\*\*,  
Toyoshige ENDŌ, and Fumio HANAOKA\*\*,

泉 雅子\*\*, 宮澤 宏\*\*, 鎌倉高志\*\*, 山口 勇\*\*,  
遠藤 豊成, 花岡文雄\*\*

Blasticidin S is a microbial antibiotic that inhibits protein synthesis in both prokaryotes and eukaryotes. The blasticidin S-resistance gene (*bsr*), isolated from *Bacillus cereus* K55-S1 strain, was inserted into pSV2 plasmid vector and introduced into cultured mammalian cells by transfection. The *bsr* gene was integrated into the genome and conferred blasticidin S resistance on HeLa cells. The transfection frequency of the *bsr* gene was as high as that of the aminoglycoside phosphotransferase gene, the so-called *neo* gene, which is a representative selectable marker for mammalian cells. Transfectants in which several copies of *bsr* had been integrated into the genome were highly resistant to blasticidin S. Furthermore, blasticidin S killed the cells more rapidly than G418, which is conventionally used as a selective drug for the *neo* gene. Thus *bsr* is concluded to be useful as a drug-resistance marker for mammalian cells.

---

\* 本報告は, Experimental Cell Research 197, 229-233 (1991) に発表

\*\* 理化学研究所