Title	New proofs of some basic theorems on stationary point processes
Sub Title	
Author	南, 就将(Minami, Nariyuki)
Publisher	慶應義塾大学日吉紀要刊行委員会
Publication year	2012
Jtitle	慶應義塾大学日吉紀要. 自然科学 (The Hiyoshi review of the natural science). No.52 (2012. 9) ,p.37- 47
JaLC DOI	
Abstract	We give new proofs of three basic theorems on stationary point processes on the real line–theorems of Khintchine, Korolyuk, and Dobrushin. Moreover we give a direct construction of the Palm measure for a class of point processes which includes stationary ones as special cases.
Notes	研究ノート
Genre	Departmental Bulletin Paper
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN10079809-20120930-0037

Hiyoshi Review of Natural Science
Keio University No. 52, 37-47 (2012)

New Proofs of Some Basic Theorems on Stationary Point Processes

Nariyuki MINAMI*

Summary——We give new proofs of three basic theorems on stationary point processes on the real line——theorems of Khintchine, Korolyuk, and Dobrushin. Moreover we give a direct construction of the Palm measure for a class of point processes which includes stationary ones as special cases.

1. Introduction

The purpose of this note is to give new proofs, based on a same simple idea, to some basic theorems on stationary point processes on the real line **R**, as stated in standard treatises on point processes such as Daley and Vere-Jones (see §3.3 of [3]).

To begin with, let us introduce necessary definitions and notation. By M_p, we denote the set of all integer-valued Radon measures on **R**. Namely M_p is the totality of all measures $N(dx)$ on **R** such that for any bounded Borel set B, $N(B)$ is a non-negative integer. Let us call any such measure a *counting measure*. For a counting measure $N \in M_p$, let us define

$$X(t) := N((0, t]) \quad (t \geq 0), \qquad := -N((t, 0)) \quad (t < 0). \tag{1}$$

Then the function $X(t)$ is right-continuous, integer-valued, locally bounded and non-decreasing. Hence $X(t)$ is piecewise constant on **R** and the set Δ, finite or countably infinite, of its points of discontinuity has no accumulation points other than $\pm\infty$. Thus the points in Δ can be ordered as

$$\cdots < x_{-1} < x_0 \leq 0 < x_1 < x_2 < \cdots,$$

so that if we let $m_n := X(x_n) - X(x_n - 0)$, then $N(dx)$ can be represented as

$$N(dx) = \sum_n m_n \delta_{x_n}(dx), \tag{2}$$

where δ_a denotes the unit mass placed at a. Each m_n is a positive integer and is called

* 南就将，慶應義塾大学医学部数学教室（〒223-8521　横浜市港北区日吉 4-1-1）：School of medicine, Keio University, Hiyoshi, Kohoku-ku, Yokohama 223-8521, Japan [Received Mar. 3, 2012]

the *multiplicity* of the point x_n. In general, either $N([0, \infty))$ or $N((-\infty, 0))$ can be finite, in which case either $\{x_n\}_{n>0}$ or $\{x_n\}_{n\le 0}$ is a finite sequence. If in the former [resp. latter] case $\{x_n\}_{n>0}$ [resp. $\{x_n\}_{n\le 0}$] terminates with x_ν, then we will set $x_n = \infty$ [resp. $x_n = -\infty$] for $n > \nu$ [resp. $n < \nu$]. When $m_n = 1$ for all n such that $x_n \neq \pm\infty$, the counting measure N is said to be *simple*. For each $N \in M_p$ with representation (2), let us associate a simple counting measure N^* defined by

$$N^*(dx) = \sum_n \delta_{x_n}(dx). \tag{3}$$

In order to make M_p a measurable space, we define $\mathcal{M}_p$ to be the σ-algebra of subsets of M_p generated by all mappings of the form

$$M_p \ni N \mapsto N(B) \in [0, \infty] \tag{4}$$

for all Borel sets $B \subset \mathbf{R}$. Then we see that x_n, m_n and N^* are all measurable functions of N, as the following lemma shows.

Lemma 1 *(i) The set*

$$C := \{N \in M_p : N((-\infty, 0]) = N((0, \infty)) = \infty\} = \{N \in M_p : x_n \textit{ is finite for all } n\}$$

belongs to $\mathcal{M}_p$.

(ii) For each integer n, x_n *and* m_n *are* $\mathcal{M}_p$*-measurable functions of N.*

(iii) The mapping $M_p \ni N \mapsto N^* \in M_p$ *is* $\mathcal{M}_p/\mathcal{M}_p$*-measurable.*

Proof. (i) The assertion is obvious from the definition of $\mathcal{M}_p$, since we can write

$$C = \bigcap_{k=1}^{\infty} \bigcup_{n=1}^{\infty} \{N \in M_p : N((-n, 0]) > k,\ N((0, n]) > k\}.$$

(ii) The measurability of x_1 follows from the relation

$$\{N \in M_p : x_1 > t\} = \{N \in M_p : N((0, t]) = 0\},$$

which holds for all $t \ge 0$. Now for each $k \ge 1$, define

$$x_1^{(k)} := \sum_{j=1}^{\infty} \frac{j}{2^n} \mathbf{1}_{((j-1)/2^n,\, j/2^n]}(x_1) + \infty \cdot 1_{\{x_1 = \infty\}}.$$

Then we see that $x_1^{(k)}$ is measurable in N and that $x_1^{(k)} \searrow x_1$ as $k \to \infty$. By the right-continuity of $X(t) = N((0, t])$ at $t > 0$, we have, as $k \to \infty$,

$$\mathbf{1}_{\{x_1 < \infty\}} \cdot X(x_1^{(k)}) = \sum_{j=1}^{\infty} \mathbf{1}_{((j-1)/2^n,\, j/2^n]}(x_1) X\left(\frac{j}{2^k}\right) \longrightarrow X(x_1) = m_1,$$

which shows the measurability of m_1 in N.

Next let $\tilde{X}(t) := X(t) - X(t \wedge x_1)$. This is measurable in N for all $t \ge 0$, since

$$X(t \wedge x_1) = X(t)\mathbf{1}_{\{x_1 \geq t\}} + X(x_1)\mathbf{1}_{\{x_1 < t\}}.$$

If we apply the above argument to $\tilde{X}(t)$ instead of $X(t)$, we can verify the measurability of x_2 and m_2 in N, and the argument can be iterated to give the measurability of all x_n and m_n.

(iii) For each $j = 0, 1, 2, \ldots$ and $t > 0$, the sets

$$\{N \in M_p : N^*((0, t]) = j\} = \{N \in M_p : x_j \leq t < x_{j+1}\}$$

and

$$\{N \in M_p : N^*((-t, 0]) = j\} = \{N \in M_p : x_{-j} \leq t < x_{-j+1}\}$$

belong to M_p. Now for each $n \geq 1$, let $\mathcal{G}_n$ be the class of all Borel subsets B of $[-n, n]$ such that the mapping

$$M_p \ni N \mapsto N^*(B) \in [0, \infty) \tag{5}$$

is measurable. Then $\mathcal{G}_n$ is seen to be a λ-system which contains the class of intervals

$$\mathcal{I} := \{(0, t] : 0 < t \leq t\} \cup \{(-t, 0] : 0 < t \leq n\}$$

which forms a π-system. Hence by Dynkin's π-λ theorem (see e.g. Durrett [2]), $\mathcal{G}_n$ contains all Borel subsets of $[-n, n]$. Since $n \geq 1$ is arbitrary, and since we can write $N^*(B) = \lim_{n\to\infty} N^*(B \cap [-n, n])$, the mapping (5) is measurable for all Borel subsets of $\mathbf{R}$.

Remark 1. By an argument similar to (iii), it is easy to show that M_p is generated by mappings $M_p \ni N \mapsto X(t)$ for all t, where $X(t)$ is defined in (1).

Definition 1 *A* point process N_ω *is a random variable defined on a probability space* $(\Omega, \mathcal{F}, \mathbf{P})$ *and taking values in the measurable space* $(M_p, \mathcal{M}_p)$.

Definition 2 *A point process* N_ω *is said to be* crudely stationary *if for any bounded interval* I *and for any* $x \in \mathbf{R}$, $N_\omega(I)$ *and* $N_\omega(I + x)$ *are identically distributed. Its* mean density *is the expectation value* $m := \mathbf{E}[N_\omega((0, 1])] \leq \infty$.

Definition 3 *A point process* N_ω *is said to be* stationary *if for any* $C \in \mathcal{M}_p$ *and* $x \in \mathbf{R}$, *one has the identity*

$$\mathbf{P}(N_\omega(\cdot) \in C) = \mathbf{P}(N_\omega(x + \cdot) \in C).$$

Obviously, N_ω is crudely stationary if it is stationary.

Remark 2. By another application of π-λ theorem, one can show without difficulty that N_ω is stationary if and only if for any finite family of Borel subsets $B_1, \ldots, B_n$ of $\mathbf{R}$, and of non-negative integers $k_1, \ldots, k_n$, the identity

$$\mathbf{P}(N_\omega(B_i) = k_i,\ i = 1, \ldots, n) = \mathbf{P}(N_\omega(x + B_i) = k_i,\ i = 1, \ldots, n)$$

holds for any $x \in \mathbf{R}$.

2. Basic theorems and their proofs

Our argument is based on the following lemma, which is an immediate consequence of Definition 2.

Lemma 2 *Let the point process N_ω be crudely stationary. Then for any bounded interval I and for any non-negative integer k,*

$$\mathbf{P}(N_\omega(I) = k) = \int_0^1 \mathbf{P}(N_\omega(x + I) = k)\,dx = \mathbf{E}\left[\int_0^1 \mathbf{1}_{\{N_\omega(x+I)=k\}}\,dx\right].$$

Proposition 1 (Khintchine's theorem) *For any crudely stationary point process N_ω, the limit*

$$\lambda := \lim_{h \searrow 0} \frac{1}{h}\mathbf{P}(N_\omega((0, h]) > 0)$$

exists and satisfies $\lambda \le m$. λ is called the intensity *of the point process N_ω.*

Proof. Let N_ω be represented as (2) and define the point process N_ω^* by (3). If we set $\nu(\omega) := N_\omega^*(0, 1]$, it satisfies $x_{\nu(\omega)}(\omega) \le 1 < x_{\nu(\omega)+1}(\omega)$. Obviously we have

$$\begin{aligned}\{x \in (0,1] : N_\omega((x, x+h]) > 0\} &= (0,1] \cap \left[\bigcup_{j=1}^{\infty}[x_j(\omega) - h, x_j(\omega))\right] \\ = (0,1] \cap \left[\bigcup_{j=1}^{\nu(\omega)+1} J_j^\omega(h)\right] &= \sum_{j=1}^{\nu(\omega)+1}\left[(0,1] \cap (J_j^\omega(h) \setminus J_{j-1}^\omega(h))\right],\end{aligned}$$

where we have set $J_j^\omega(h) := [x_j(\omega) - h, x_j(\omega))$ and $J_0 = \emptyset$. Hence

$$\begin{aligned}\frac{1}{h}\int_0^1 \mathbf{1}_{\{N_\omega((x,x+h])>0\}}\,dx &= \frac{1}{h}\sum_{j=1}^{\nu(\omega)+1} |(0,1] \cap (J_j^\omega(\mathrm{h}) \setminus J_{j-1}^\omega(h))| \\ &= \sum_{j=1}^{\nu(\omega)+1} \frac{1}{h}\{(1 \wedge x_j(\omega)) - (0 \vee x_{j-1}(\omega) \vee (x_j(\omega) - h)\}_+ ,\end{aligned}$$

where for a Borel subset B of $\mathbf{R}$, $|B|$ denotes its Lebesgue measure and for a real number a, $a_+ := a \vee 0 = \max\{a, 0\}$ denotes its positive part. Now it is easy to see that for $1 \le j \le \nu(\omega)$,

$$\frac{1}{h}\{(1 \wedge x_j(\omega)) - (0 \vee x_{j-1}(\omega) \vee (x_j(\omega) - h)\}_+ \nearrow 1$$

as $h \searrow 0$, and that for $j = \nu(\omega) + 1$,

$$\frac{1}{h}\{(1 \wedge x_{\nu(\omega)+1}(\omega)) - (0 \vee x_{\nu(\omega)}(\omega) \vee (x_{\nu(\omega)+1}(\omega) - h))\}_+$$

is bounded by 1 and tends to 0 as $h \searrow 0$. Thus we can apply the monotone convergence theorem, the dominated convergence theorem and Lemma 2, to obtain

$$\begin{aligned} \frac{1}{h}\mathbf{P}(N_\omega((0, h]) > 0) &= \mathrm{E}\left[\frac{1}{h}\sum_{j=1}^{\nu(\omega)+1} |(0, 1] \cap (J_j^\omega \setminus J_{j-1}^\omega(h))|\right] \\ &\to \mathrm{E}\left[\sum_{j=1}^{\nu(\omega)} 1\right] = \mathrm{E}\left[N_\omega^*((0, 1])\right], \end{aligned}$$

as $h \searrow 0$. Thus the desired limit λ exists and is equal to $\mathbf{E}[N_\omega^*((0, 1])]$. Clearly it satisfies the inequality $\lambda \le \mathbf{E}[N_\omega((0, 1])] = m$.

Corollary 1 *If N_ω is simple, then $\lambda = m$. When $m < \infty$, the converse is also true.*

Proof. N_ω is simple if and only if $N_\omega^* = N_\omega$ almost surely, which obviously implies $\lambda = m$. On the other hand, if $\lambda = m < \infty$, then

$$\mathbf{E}[N_\omega((0, 1]) - N_\omega^*((0, 1])] = m - \lambda = 0.$$

But $N_\omega((0, 1]) - N_\omega^*((0, 1]) \ge 0$ in general, so that $N_\omega((0, 1]) = N_\omega^*((0, 1])$ almost surely. The same argument is valid if the interval $(0, 1]$ is replaced by $(n, n + 1]$, so that $N_\omega((n, n + 1]) = N_\omega^*((n, n + 1])$ almost surely for all integers n, and the simplicity of N_ω follows.

Remark 3. In the treatise by Daley and Vere-Jones [3], for example, Proposition 1 is proved in the following way: If we define $\varphi(h) := \mathbf{P}(N_\omega((0, h]) > 0)$, then by the crude stationarity, we have for any positive h_1 and h_2,

$$\begin{aligned} \varphi(h_1 + h_2) &= \mathbf{P}(N_\omega((0, h_1 + h_2]) > 0) = \mathbf{P}(N_\omega((0, h_1]) + N_\omega((h_1, h_1 + h_2]) > 0) \\ &\le \mathbf{P}(N_\omega((0, h_1]) > 0) + \mathbf{P}(N_\omega((h_1, h_1 + h_2]) > 0) = \varphi(h_1) + \varphi(h_2), \end{aligned}$$

so that $\varphi(h)$ is a sub-additive function defined on $[0, \infty)$ satisfying $\varphi(0) = 0$. To show the existence of the intensity λ, it suffices to apply the following well known lemma.

Lemma 3 *Let $g(x)$ be a sub-additive function defined on $[0, \infty)$ such that $g(0) = 0$. Then one has*

$$\lim_{x \searrow 0} \frac{g(x)}{x} = \sup_{x > 0} \frac{g(x)}{x} \leq \infty .$$

However, this argument does not provide the representation $\lambda = \mathbf{E}[N^*_\omega((0, 1])]$, so that the proof of Corollary 1 requires some extra work. Our proof above is closer to that of Leadbetter [5]. See also Chung [1].

Definition 4 *A crudely stationary point process N_ω is said to be* orderly *when*

$$\mathbf{P}(N_\omega((0, h]) \geq 2) = o(h) \quad (h \searrow 0).$$

Proposition 2 (Dobrushin's theorem) *If a crudely stationary point process N_ω is simple and if $\lambda < \infty$, then N_ω is orderly.*

Proof. By Lemma 2, we can write

$$\mathbf{P}(N_\omega((0, h]) \geq 2) = \mathbf{E}\left[\int_0^1 \mathbf{1}_{\{N_\omega((x, x+h]) \geq 2\}} dx\right].$$

As can be seen from the proof of Proposition 1, we have

$$\begin{aligned} & \frac{1}{h}\int_0^1 \mathbf{1}_{\{N_\omega((x,x+h]) \geq 2\}} dx \leq \frac{1}{h}\int_0^1 \mathbf{1}_{\{N_\omega((x,x+h]) > 0\}} dx \\ = & \sum_{j=1}^{\nu(\omega)} \frac{1}{h} |(0, 1] \cap (J_j^\omega(h) \setminus J_{j-1}^\omega(h))| + \frac{1}{h} |(0, 1] \cap (J_{\nu(\omega)+1}(h) \setminus J_{\nu(\omega)}(h)| \\ \leq & N^*_\omega((0, 1]) + 1 , \end{aligned}$$

and

$$\lim_{h \searrow 0} \frac{1}{h}\int_0^1 \mathbf{1}_{\{N_\omega((x,x+h]) \geq 2\}} dx = \sharp\{j : x_j(\omega) \in (0, 1],\ m_j(\omega) \geq 2\} .$$

Since $\mathbf{E}[N^*_\omega((0, 1])] = \lambda < \infty$, we can apply the dominated convergence theorem, to obtain

$$\lim_{h \searrow 0} \frac{1}{h}\mathbf{P}(N_\omega((0, h]) \geq 2) = \mathbf{E}[\sharp\{j : x_j(\omega) \in (0, 1],\ m_j(\omega) \geq 2\}],$$

which is equal to 0 if N_ω is simple.

Remark 4. The condition $\lambda < \infty$ cannot be dropped. For a counter example, see Exercise 3.3.2 of [3].

Proposition 3 (Korolyuk's theorem) *A crudely stationary, orderly point process is simple.*

Proof. By Fatou's lemma and the orderliness of N_ω,

$$\begin{aligned}\mathbf{E}[\sharp\{j : x_j(\omega) \in (0,1],\ m_j(\omega) \geq 2\}] &= \mathbf{E}\left[\liminf_{h\searrow 0}\frac{1}{h}\int_0^1 \mathbf{1}_{\{N_\omega((x,x+h])\geq 2\}}\,dx\right] \\ &\leq \liminf_{h\searrow 0}\frac{1}{h}\mathbf{P}(N_\omega((0,h]) \geq 2) = 0\ ,\end{aligned}$$

so that with probability one, N_ω has no multiple points in (0, 1]. By crude stationarity, the above argument is also valid if (0, 1] is replaced by (n, n+1] for any integer n. Hence N_ω is simple.

Proposition 4 *For a crudely stationary point process N_ω with finite intensity λ, the limits*

$$\lambda_k := \lim_{h\searrow 0}\frac{1}{h}\mathbf{P}(1 \leq N_\omega((0,h]) \leq k)$$

exists for $k = 1, 2, \ldots$, and satisfy $\lambda_k \nearrow \lambda$ as $k \to \infty$. Moreover for $k = 1, 2, \ldots$,

$$\pi_k := \frac{\lambda_k - \lambda_{k-1}}{\lambda} = \lim_{h\searrow 0}\mathbf{P}(N_\omega((0,h]) = k \mid N_\omega((0,h]) > 0)\ ,$$

where we set $\lambda_0 := 0$.

Proof. As before, one has

$$\frac{1}{h}\int_0^1 \mathbf{1}_{\{1\leq N_\omega((x,x+h])\leq k\}}\,dx \leq 1 + N^*_\omega((0,1])\ ,$$

and

$$\lim_{h\searrow 0}\frac{1}{h}\int_0^1 \mathbf{1}_{\{1\leq N_\omega((x,x+h])\leq k\}}\,dx = \sharp\{j : x_j(\omega) \in (0,1],\ m_j(\omega) \leq k\}\ .$$

Since $\lambda = \mathbf{E}[N_\omega^*((0,1])] < \infty$, we can apply the dominated convergence theorem and Lemma 2, to obtain

$$\lambda_k = \lim_{h\searrow 0}\frac{1}{h}\mathbf{E}\left[\int_0^1 \mathbf{1}_{\{1\leq N_\omega((x,x+h])\leq k\}}\,dx\right] = \mathbf{E}[\sharp\{j : x_j(\omega) \in (0,1],\ m_j(\omega) \leq k\}]\ .$$

This representation of λ_k immediately gives

$$\lim_{k\to\infty}\lambda_k = \mathbf{E}[\sharp\{j : x_j(\omega) \in (0,1]\}] = \mathbf{E}[N^*((0,1])] = \lambda,$$

by the monotone convergence theorem. The last statement of the proposition is obvious.

Corollary 2 *For a crudely stationary point process with finite intensity, we have*

$$\lambda\sum_{k=1}^{\infty} k\pi_k = \mathbf{E}[N_\omega((0,1])] = m\ .$$

3. The Palm measure

Let us assume that the probability space $(\Omega, \mathcal{F}, \mathbf{P})$, on which our point process N_ω is defined, is equipped with a measurable flow $\{\theta_t\}_{t \in \mathbf{R}}$. Here a measurable flow $\{\theta_t\}$ is, by definition, a family of bijections $\theta_t : \Omega \to \Omega$ such that

(a) θ_0 is the identity mapping, and for any $s, t \in \mathbf{R}$, $\theta_s \circ \theta_t = \theta_{s+t}$ holds;
(b) the mapping $(t, \omega) \mapsto \theta_t(\omega)$ from $\mathbf{R} \times \Omega$ into Ω is jointly measurable with respect to $\mathcal{B}(\mathbf{R}) \times \mathcal{F}$, where $\mathcal{B}(\mathbf{R})$ is the Borel σ-algebra on $\mathbf{R}$.

Let us further assume that the relation

$$\int_{\mathbf{R}} N_{\theta_t \omega}(dx)\varphi(x) = \int_{\mathbf{R}} N_\omega(dx)\varphi(x - t) \tag{6}$$

holds for any $t \in \mathbf{R}$ and any continuous function φ with compact support. If the probability measure $\mathbf{P}$ is $\{\theta_t\}$-invariant in the sense $\mathbf{P} \circ \theta_t^{-1} = \mathbf{P}$ for all $t \in \mathbf{R}$, then by (6), our point process N_ω is stationary.

Definition 5 *The Palm measure of a point process $N_\omega(dx)$ is a measure kernel $Q(x, d\omega)$ on $\mathbf{R} \times \Omega$ such that for any jointly measurable, non-negative function $f(x, \omega)$, the relation*

$$\int_\Omega \mathbf{P}(d\omega) \int_{\mathbf{R}} N_\omega(dx) f(x, \omega) = \int_{\mathbf{R}} \lambda(dx) \int_\Omega Q(x, d\omega) f(x, \omega) \tag{7}$$

holds, where $\lambda(dx)$ is the mean measure of N_ω which is defined by $\lambda(B) = \mathrm{E}[N_\omega(B)]$ for $B \in \mathcal{B}(\mathbf{R})$ and which we assume to be finite for bounded Borel sets B.

Now let $u(t)$ be a probability density function on $\mathbf{R}$. Define a new probability measure $\mathbf{P}_u$ by

$$\int_\Omega \mathbf{P}_u(d\omega) g(\omega) = \int_{\mathbf{R}} u(t)\, dt \left(\int_\Omega \mathbf{P}(d\omega) g(\theta_t \omega) \right), \tag{8}$$

where $g(\omega)$ is an arbitrary non-negative measurable function on Ω. Then the following result holds.

Theorem 1 *For any probability density $u(t)$ on $\mathbf{R}$, the Palm measure $Q_u(x, d\omega)$ exists for the point process N_ω defined on the probability space $(\Omega, \mathcal{F}, \mathbf{P}_u)$.*

Proof. Let $f(x, \omega) \geq 0$ be jointly measurable on $\mathbf{R} \times \Omega$. Then we can rewrite the left

hand side of (7) in the following way:

$$\begin{aligned}\int_\Omega \mathbf{P}_u(d\omega)\int_{\mathbf{R}} N_\omega(dx)f(x,\,\omega) &= \int_{\mathbf{R}} u(t)dt\int_\Omega \mathbf{P}(d\omega)\int_{\mathbf{R}} N_{\theta_t\omega}(dx)f(x,\,\theta_t\omega)\\ &= \int_{\mathbf{R}} u(t)dt\int_\Omega \mathbf{P}(d\omega)\int_{\mathbf{R}} \mathrm{N}_\omega(dx)f(x-t,\,\theta_t\omega)\\ &= \int_\Omega \mathbf{P}(d\omega)\int_{\mathbf{R}} N_\omega(dx)\int_{\mathbf{R}} u(t)dtf(x-t,\;\theta_t\,\omega)\\ &= \int_\Omega \mathbf{P}(d\omega)\int_{\mathbf{R}} N_\omega(dx)\int_{\mathbf{R}} u(x-s)dsf(\mathrm{s},\,\theta_{x-s}\,\omega)\\ &= \int_{\mathbf{R}} ds\int_\Omega \mathbf{P}(d\omega)\int_{\mathbf{R}} N_\omega(dx)u(x-s)f(s,\,\theta_{x-s}\,\omega)\,. \qquad (9)\end{aligned}$$

At this stage, take $f(x,\;\omega) = \varphi(x)$. Then (9) reduces to

$$\int_{\mathbf{R}} \varphi(s)\lambda(ds) = \int_{\mathbf{R}} \varphi(s)\ell_u(s)ds \qquad (10)$$

with

$$\ell_u(s) = \int_\Omega \mathbf{P}(d\omega)\int_{\mathbf{R}} N_\omega(dx)u(x-s)\ . \qquad (11)$$

If we define, for each $s \in \mathbf{R}$, the measure $Q_u(s,\,d\omega)$ on $(\Omega,\,\mathcal{F})$ by

$$\int_\Omega Q_u(s,d\omega)\,g(\omega) = \frac{\mathbf{1}_{(0,\infty)}(\ell_u(s))}{\ell_u(s)}\int_\Omega \mathbf{P}(d\omega)\int_{\mathbf{R}} N_\omega(dx)\,u(x-s)\,g(\theta_{x-s}\omega)\ , \qquad (12)$$

then (9) takes the form of (7), and the theorem is proved.

When $\mathbf{P}$ is $\{\theta_t\}$-invariant, then we have $\mathbf{P}_u = \mathbf{P}$ for any probability density u on $\mathbf{R}$, and

$$\ell_u(s) = \int_\Omega \mathbf{P}(d\omega)\int_{\mathbf{R}} N_{\theta_s\omega}(dx)\,u(x) = \int_\Omega \mathbf{P}(d\omega)\int_{\mathbf{R}} N_\omega(dx)\,u(x) =: \ell > 0$$

is a constant. Moreover one can compute as

$$\begin{aligned}&\int_\Omega Q_u(s,d\omega)\,g(\omega) = \frac{1}{\ell}\int_\Omega \mathbf{P}(d\omega)\int_{\mathbf{R}} N_{\theta_s\omega}(dx)u(x)\,g(\theta_x\,\omega)\\ =\ &\frac{1}{\ell}\int_\Omega \mathbf{P}(d\omega)\int_{\mathbf{R}} N_{\theta_s\omega}(dx)\,u(x)\,g(\theta_{x-s}(\theta_s\omega)) = \frac{1}{\ell}\int_\Omega \mathbf{P}(d\omega)\int_{\mathbf{R}} N_\omega(dx)\,u(x)\,g(\theta_{x-s}\omega)\ .\end{aligned}$$

Hence if we define a measure $\hat{\mathbf{P}}\,(d\omega)$ on $(\Omega,\,\mathcal{F})$ by

$$\int_\Omega \hat{\mathbf{P}}(d\omega)g(\omega) = \int_\Omega \mathbf{P}(d\omega)\int_{\mathbf{R}} N_\omega(dx)\,u(x)\,g(\theta_x\,\omega)\ ,$$

then we get

$$Q_u(s,d\omega) = \frac{1}{\ell}(\hat{\mathbf{P}}\circ\theta_s)(d\omega)\ ,$$

and (7) can be written in the form

$$\int_{\Omega} \mathbf{P}(d\omega) \int_{\mathbf{R}} N_{\omega}(dx) f(x, \omega) = \int_{\mathbf{R}} dx \int_{\Omega} \hat{\mathbf{P}}(d\omega) f(x, \theta_{-x}\omega) \ , \tag{13}$$

which is the defining relation of the Palm measure in the stationary case (see [6]). (13) shows in particular that the definition of $\hat{\mathbf{P}}$ is independent of the choice of u.

Our consideration of the probability measure $\mathbf{P}_u$ is motivated by the following observation.

Proposition 5 *The probability measure* $\mathbf{P}$ *is* $\{\theta_t\}$*-invariant if and only if the following two conditions hold:*

(i) $\mathbf{P}_u = \mathbf{P}$ *for any probability density function* $u(t)$ *on* $\mathbf{R}$*;*
(ii) *the set* H *of all bounded measurable functions* $\varphi(\omega)$ *on* Ω *such that* $t \mapsto \varphi(\theta_t\omega)$ *is continuous for all* $\omega \in \Omega$ *is dense in* $L^2(\Omega, \mathbf{P})$.

Proof. The necessity of (i) is obvious. That (ii) also follows from the $\{\theta_t\}$-invariance of $\mathbf{P}$ is proved in [6] (see Lemma II. 3). To prove the sufficiency of (i) and (ii), fix an arbitrary $t_0 \in \mathbf{R}$ and take a sequence of probability density $\{u_n\}_n$ so that $u_n(t)dt \to \delta_{t_0}(dt)$ weakly. Now for any $\varphi \in H$, $t \mapsto \varphi(\theta_t\omega)$ is continuous and bounded by $\|\varphi\|\infty := \sup_{\Omega} |\varphi(\omega)|$. Hence we can apply the dominated convergence theorem, to get

$$\begin{aligned}
\int_{\Omega} \mathbf{P}(d\omega)\varphi(\theta_{t_0}\omega) &= \int_{\Omega} \mathbf{P}(d\omega) \left(\lim_{n\to\infty} \int_{\mathbf{R}} \varphi(\theta_t\omega) u_n(t) dt \right) \\
&= \lim_{n\to\infty} \int_{\mathbf{R}} \left(\int_{\Omega} \mathbf{P}(d\omega)\varphi(\theta_t\omega) \right) u_n(t) dt \\
&= \lim_{n\to\infty} \int_{\Omega} \mathbf{P}_{u_n}(d\omega)\, \varphi(\omega) = \int_{\Omega} \mathbf{P}(d\omega)\, \varphi(\omega)
\end{aligned}$$

by condition (i). But if H is dense in $L^2(\Omega, \mathbf{P})$, we can approximate an arbitrary bounded measurable function $g(\omega)$ by the elements of H, to obtain

$$\int_{\Omega} \mathbf{P}(d\omega) g(\theta_{t_0}\omega) = \int_{\Omega} \mathbf{P}(d\omega) g(\omega)$$

for any $t_0 \in \mathbf{R}$. This sows the $\{\theta_t\}$-invariance of $\mathbf{P}$.

In most cases of application, Ω itself is a topological space with $\mathcal{F}$ the Baire σ-algebra generated by that topology and $t \mapsto \theta_t\omega$ is continuous for all $\omega \in \Omega$. In such a case, H contains the class $C_b(\Omega)$ of all bounded continuous functions on Ω, which is dense in $L^2(\Omega, \mathbf{P})$. Hence condition (ii) is not as restrictive as it may appear.

See [4] for a general treatment of stationary random measures on a topological group.

References

[1] Chung, K. L.: Crudely stationary point processes, *Amer. Math. Monthly* **79**, (1972) 867–877.

[2] Durrett, R.: *Probability: theory and examples*, Second edition, Duxbury Press, Belmont, CA, 1996.

[3] Daley, D.J. and Vere-Jones, D.: *An introduction to the theory of point processes. I: Elementary theory and methods*, 2nd ed., Probab. Appl. (N. Y.), Springer, New York, 2003.

[4] Last, G.: Modern random measures: Palm theory and related models, in *New perspectives in stochastic geometry* (W. S. Kendall and I. Molchanov, eds.), Oxford 2010.

[5] Leadbetter, M.R.: On three basic results in the theory of stationary point processes, *Proc. Amer. Math. Soc.* **19**, (1968) 115–117.

[6] Neveu, J.: Processus ponctuels, École d'Eté de Probabilités de Saint-Flour VI. *Lecture Notes in Mathematics* **598**, 249–445, Springer, Berlin.