Title	Spectrophotometric determination of palladium with a new chelating reagent, 1-methyl-3-(4-methyl-2-thiazolyl) thiourea
Sub Title	
Author	八木, 優子(Yagi, Yuko) 堀内, 公子(Horiuchi, Kimiko) 与田, 玲子(Yoda, Reiko) 山本, 有一(Yamamoto, Yuichi) 村上, 悠紀雄(Murakami, Yukio)
Publisher	共立薬科大学
Publication year	1975
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.20 (1975.) ,p.123- 123
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000020-0123

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって 保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Spectorophotometric Determination of Palladium with a New Chelating Reagent, 1-Methyl-3-(4-methyl-2-thiazolyl) thiourea*

YUKO YAGI, KIMIKO HORIUCHI, REIKO YODA, YUICHI YAMAMOTO and YUKIO MURAKAMI

八木 優子, 堀内 公子, 与田 玲子 山本 有一, 村上悠紀雄

Out of 47 thioureid derivatives synthesized, the title compound was chosen to perform a screenting test on the chelate formation with 66 metal ions. The formation of colored chelate was visually observed for Pd^{2+} , Cu^{2+} , Pt^{4+} , Au^{3+} , Ru^{3+} and Ag^+ in acidic media. The highest sensitivity was obtained for Pd chelate and the chelate formation was completed within five minutes at PH 2 at the room temperature. Conformity to Beer's Law is sufficiently good up to 4 ppm of Pd. The Pd chelate was readily extracted with chloroform and exhibited the maximum absorption at 280 and 320 nm. The excess reagent, causing an absorption peak at 290 nm, could be removed by shaking with 1 mol/1 NaOH solution, while the chelate remained unchanged. The molar absorption coefficient was 1.76×10^4 (reference : H_2O) at 320 nm and 2.40×10^4 (reference : $CHCl_3$) at 280 nm. According to Ringbom's plot, the optimum concentration range was 0.8-4.4 ppm for the 320 nm peak and 0.6-3.2 ppm for 280 nm peak. Pretreatment of sample solution with perchloric acid was effective to reduce the interference of Cl^- , OsO_4 or Ru^{3+} and also to reduce the aging effect of the Pd solution of $10^{-3}-10^{-4}$ mol/1.

The proposed procedure is as follows. Add several drops of perchloric acid to the sample solution containing up to 40 μ g Pd and heat to almost dryness. Make up the solution nearly 10 ml and adjust the pH to about 2 with 1 mol/1 NaOH solution. After adding 1 ml of the reagent solution (2×10⁻³mol/1), extract the chelate by shaking with 8 ml of chloroform. Allow to stand for 5 minutes to fully separate the two layers. Then scrub the organic layer to remove the excess reagent with equal volume of 1 mol/1 NaOH solution. Measure the absorbance at 280 nm in the chloroform as reference solution.

^{*}本報告は、日本化学会誌 5,831-835,(1975)に発表.