Title	Incorporation of [2-³H]- and [2-²H₃]-acetate by penicillium urticae into griseofulvin and determination of the stereochemistry of isotopes at C-5'			
Sub Title				
Author	佐藤, 良博(Sato, Yoshihiro) 町田(関), 敏子(Machida(Seki), Toshiko) 小田, 泰子(Oda, Taiko)			
Publisher	共立薬科大学			
Publication year	1976			
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.21 (1976.) ,p.96- 98			
JaLC DOI				
Abstract				
Notes	抄録			
Genre	Technical Report			
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000021-0096			

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって 保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

INCORPORATION OF (2-³H)- AND (2-²H₃)-ACETATE BY PENICILLIUM URTICAE INTO GRISEOFULVIN AND DETERMINATION OF THE STEREOCHEMISTRY OF ISOTOPES AT C-5' *

Yoshihiro Sato, Toshiko Machida (Seki) and Taiko Oda 佐藤良博,町田(関)敏子,小田泰子

Griseofulvin is known as a typical example of polyketomethylene antibiotics. The biosynthetic pathway of griseofulvin from acetate-malonate units has already been established by Birch's and Rhode's groups using ¹⁴C or ³⁶Cl-precursors. However, problems regarding the extent of incorporation of tritiums or deuteriums from $(2^{-3}H)$ - or $(2^{-2}H_3)$ -acetate into griseofulvin have remained to be solved.

In this communication we wish to report the incorporation of tritium from $[2^{-3}H]$ -acetate into the griseofulvin molecule by *Penicillium urticae*. The labelled precursor consisting of 1.08×10^7 dpm of $[2^{-14}C]$ -acetate and 6.77×10^7 dpm of $[2^{-3}H]$ acetate $(^{3}H)^{14}C$ ratio, 6.27) was added to the shaking culture of *P. urticae* on the 8 th day of the fermentation period. After 2 or 24 hr of fermentation, broth from each flask was filtered to separate the mycelium and filtrate, both of which were subsequently separated into neutral and acidic fractions. Griseofulvin was then obtained from the neutral fraction. The result is shown in Table I.

Fermentation time (hr)	Isotope	dpm/mg	% of Incorp.	³H∕¹ 4C
0	³Н	7.99×10^{3}	0.46	1.85
2	1 4 C	4.32×10^{3}	1.43	
24	³Н	1. 29 × 104	0. 55	1. 77
	1 4 C	7.27×10^3	1.78	

TABLE I. Radioactivity of Griseofulvin obtained from the Filtrate

The results indicate that, during the biosynthesis of griseofulvin through polyketomethylene intermediate (II), all the tritium atoms on griseophenone B (III) were retained in the griseofulvin molecule (Chart 1).

The positions of tritium atoms incorporated into griseofulvin were proved by chemical means : Hydrolysis of the doubly labelled griseofulvin (${}^{3}H/{}^{14}C$ ratio, 2.97) with H₂SO₄ in aceti cacid gave griseofulvic acid (IV), whose ${}^{3}H/{}^{14}C$ ratio was reduced to 2.56. This shows that one tritium atom is lost from position 3'. Adsorption of a

^{*} 本報告は Tetrahedron Letters No. 51, pp4571-4574 (1975) に発表

No. 21 (1976)



Chart 1. Biosynthesis of Griseofulvin from [2-³H]-Acetate (Tritium is abbreviated as T in the chart)

chloroform solution of griseofulvin (${}^{3}H/{}^{14}C$ ratio, 2.77) on neutral alumina (Woelm, activity II) overnight afforded 5,3'-dichlorogriseofulvin (V), whose radioactivity ratio (1.87) corresponds to the lose of tritiums from positions 5 and 3'. In other experiment, a mixture of griseofulvin and neutral alumina was stirred in chloroform for 16 hr according to the Levine's method. This procedure is known to effect stereoselective partial replacement of 5' β -deuterium of griseofulvin-5', 5'-d₂ by hydrogen. The product actaully obtained by this experiment, however, showed no change of radioactivity ratio, excluding the possibility of β -orientation of tritium at position 5'. Next, griseofulvin (${}^{3}H/{}^{14}C$ ratio, 2.57) was heated under reflux for 16 hr in CH₃CH₂-CH₂C(CH₃)₂OH containing CH₃CH₂CH₂C(CH₃)₂ONa, conditions known to effect some exchange of both of the C-5' protons. The product (VI, ${}^{3}H/{}^{14}C$ ratio, 2.45) indicated a loss of radioactivity corresponding to about 30% of one tritium atom (Chart 2). This considerably small value (30%) was proved to be effective for α -orientation of tritium at C-5', because the deuteration rates at 5' α of griseofulvin in parallel ex-



- 97 -

No. 21 (1976)

periments were about 30-50% of theoretical value on ¹H nmr and mass spectrometric analyses.

In a separate experiment $[2^{-2}H_3]$ -acetate was fermented with the mycelium of *P. urticae* in the medium consisting of 5.0g of KCl, 0.5g of MgSO₄ • 7H₂O, 15.0g of sodium $[2^{-2}H_3]$ -acetate • 3H₂O (obtained from acetic acid-d₄, 99.5 Atom% D) and H₂O to a total volume of 11 (pH 7.9). An isotopic composition of $[^2H]$ -griseofulvin thus obtained was proved mass spectrometrically to consist of a deuterated mixture: d₀ 61.4, d₁ 4.3, d₂ 5.9, d₈ 8.6, d₄ 10.6, d₅ 6.4, d₆ 1.4, and d₇ and d₈ < 1%. Using CH₃O sinal as a standard, ¹H nmr spectrum was examined for the decrease in each ¹H signal intensity. The positions of tritium or deuterium atoms incorporated into griseofulvin are 5 in the A ring, and 3', 5' α , and 6'-methyl in the C ring. This result indicates the *trans* diaxial type reduction on C ring during the course of griseofulvin biosynthesis.