Title	Production of ⁶¹ Cu by α and ³ He bombardment on cobalt targets
Sub Title	
Author	本間, 義夫(Honma, Yoshio)
	村上, 悠紀雄(Murakami, Yukio)
Publisher	共立薬科大学
Publication year	1977
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of
	Pharmacy). No.22 (1977.) ,p.127- 128
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000022-0127

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって 保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Production of ⁶¹Cu by α and ³He Bombardment on Cobalt Targets^{*}

YOSHIO HOMMA and YUKIO MURAKAMI

本間義夫, 村上悠紀雄

⁶¹Cu has better nuclear properties for use in nuclear medicine.¹⁾ Its 3.32h half-life and 284 keV γ -ray make it a particularly useful diagnostic scanning agent giving a much lower absorbed dose for a given count rate than the more readily available ⁶⁴Cu and ⁶⁷Cu. ⁶¹Cu can be produced with a cyclotron by a ⁵⁹Co(α , 2n) ⁶¹Cu reaction : this reaction results in a maximum of 6 mCi/ μ Ah for 40 MeV α bombardment.¹⁾ Bombardment of cobalt has an additional advantage in that the inexpensive monoisotopic element cobalt can be used.

In producing a short-lived radionuclide for use in clinical diagnostic procedures, two factors of prime importance are the yield of the desired nuclide and the degree of contamination with other isotopes, particularly those which have relatively long half-lives and which cannot be separated chemically. To determine the optimum irradiation condition to maximize the yield of the desired nuclide and to minimize the yield of other by-product nuclides, the excitation curves for the reactions concerned must be known.

We have investigated the excitation curves and the thick-target yield curves of ${}^{61}Cu$ and by-product nuclides such as ${}^{56}Co$, ${}^{57}Co$, and ${}^{58}Co$. Photopeak efficiencies have been calculated at γ -ray energies of these nuclides for sodium iodide crystals.

Target Preparation. A thin cobalt target (15–25 mg/cm²) was electro-deposited from a cobalt sulfate solution (CoSO₄·7H₂O 500g/1, NaCl 17g/1, H₃BO₄ 45g/1) onto a disk of electrolytic copper foil (35 μ thick).

Bombardment. The stacked target was attached to the beam duct of No. 2 of IPCR cyclotron and bombarded with $0.5-1 \mu A$ beam of 40 MeV α and ³He particles. A collimater situated in front of the target reduced the spead in width of beam to ca. 1.5×1.5 cm². The beam current was measured with a beam current integrator.

Measurement. After bombardment, γ -ray spectra of each foil were measured with a 15cm³ Ge(Li) detector, which had been accurately calibrated using IAEA γ -ray standard sources. The specific γ -rays and half-lives were sufficiently distinguished without chemical separation. The principal photopeaks of nuclide were followed in order to determine the half-life and confirm the identily of the nuclides.

The excitation curves for the production of ${}^{61}Cu$ by α bombardment on cobalt are shown in Fig. 1. The maximum cross section of 350mb is shown at 25 MeV. Above 25MeV the cross section for ${}^{61}Cu$ production decreases; probably because the ${}^{59}Co(\alpha, \alpha n){}^{58}Co$ and ${}^{59}Co(\alpha, \alpha 2n){}^{57}Co$ reactions are more probable than the ${}^{59}Co(\alpha, 2n){}^{61}Cu$ reaction in this region.

^{*} 本報告は Bull. Chem. Soc. Japan, 50, 1251-1255 (1977) に発表

¹⁾ Y. Homma and Y. Murakami, Chem. Lett, 1976, 397.

No. 22 (1977)

The Coulomb barrier for the interaction of ³He with ⁵⁹Co is about 9.72MeV, whereas the Q values for the ⁵⁹Co(³He,n)⁶¹Cu, ⁵⁹Co(³He, α)⁵⁸Co, ⁵⁹Co(³He, α n)⁵⁷Co, and ⁵⁹Co(³He, α 2n)⁵⁶Co reactions are +6.6, +10.1, +1.5, and -9.4 MeV, respectively. This indicates that, for the ³He particles with sufficient kinetic energy to cross the Coulomb barrier, the cross section for the first reaction is negligibly small. (Fig. 2).

We found the bombardment of cobalt at 40 MeV to be the best method to produce ⁶¹Cu in sufficient quantity for radiopharmaceutical studies. Advantages are the relatively high yield as compared with other methods, and the high radionuclidic purity of carrier-free ⁶¹Cu. For routine production we have chosen the following bombardment conditions : energy of the incident particles 40 MeV, target thickness 250 mg/cm², beam current 5μ A. The irradiation time varies according to the required quantity of ⁶¹Cu from 1—2h. Under these conditions, the yield of ⁶¹Cu obtained at the end of 1 and 2h of bombardment is *ca*. 13 and 26mCi, respectively, corrected for losses during the course of recovery.



Fig. I Excitation curve and thick-target yield curve for α reaction on cobalt producing ⁶¹Cu.



Fig. 2 Excitation curve and thick-target yield curve for ³He reaction on cobalt producing ⁶¹Cu.