Title	Studies on thiazoline derivatives (XV) reactions of 2-phenyl and 2-methylamino-2-thiazolines with methyl and phenylisothiocyanates
Sub Title	
Author	山本, 有一(Yamamoto, Yuichi) 与田, 玲子(Yoda, Reiko) 小口, 敬子(Koguchi, Kyoko)
Publisher	共立薬科大学
Publication year	1978
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.23 (1978.) ,p.38- 46
JaLC DOI	
Abstract	
Notes	原報
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000023- 0038

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって 保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Studies on Thiazoline Derivatives (XV) Reactions of 2-Phenyl and 2-Methylamino-2-thiazolines with Methyl and Phenylisothiocyanates

YUICHI YAMAMOTO, REIKO YODA, and KYOKO KOGUCHI

山本有一, 与田玲子, 小口敬子

(Received September 30, 1978)

Summary

As a secondary amine, 2-methylamino-2-thiazoline (1b) or 2-phenylamino-2-thiazoline (1c) reacted with methylisothiocyanate (2b) or phenylisothiocyanate (2c), as shown in Scheme I, regardless of R, R'=alkyl or aryl to give 3-(N-methyl-(or phenyl)thiocarbamoyl)-2-methyl (or phenyl)iminothiazolidine (3c, d, e), but not the compound (4). There was no thermal rearrangement of (3) to (4).

The authors¹⁾ reported previously that 2-amino-2-thiazoline (1a) reacted with methylisothiocyanate (2a) under 40°C to give 3-(N-methylthiocarbamoyl)-2-iminothiazolidine (3a) which isomerized to thermodynamically more stable 2-(N-methylthiocarbamoyl)amino-2-thiazoline (4a).

It was reported that the reaction of (1a) with phenylisothiocyanate (2b) gave 3-(N-phenylthiocarbamoyl)-2-iminothiazolidine (3c) under 50°C and (3c) thermally isomerized to 2-(N-phenylthiocarbamoyl)amino-2-thiazoline (4c).² Later it was confirmedthat only (4c) was obtained regardless of the reaction temperature.³

The authors⁴) reported earlier that a secondary amine, 2-methylamino-2-thiazoline (1b), reacted with (2a) to give only 3-(N-methylthiocarbamoyl)-2-methyliminothiazoline (3b), but not the isomeric 2-(N-methylthiocarbamoyl)methylamino-2-thiazoline (4b) and that there was no thermal isomerization of (3b) to (4b).

This report has clarified the reaction between the secondary amines (1b, c) and phenylisothiocyanate (2b). It was reported that the reaction between (1c) and (2b) gave 2-(N-phenylthiocarbamoyl)phenylamino-2-thiazoline (4c),⁵⁾ however, we have shown

Y. Yamamoto and R. Yoda, Annual Report Kyoritsu Pharm. College., 18, 53 (1973); C.A., 81, 136039z (1974)

²⁾ E. Fromm and Kapeller-Alder, Ann. Chem., 467, 259 (1928)

³⁾ D.L. Klayman, J.J. Maul and George W.A. Milne, J. Hetero. Chem., 5, 520 (1968)

⁴⁾ Y. Yamamoto, R. Yoda and M. Matsumura, Chem. Pharm. Bull. (Tokyo) 23, 2134 (1975)

⁵⁾ E. Cherbuliez, Br. Baehler, S. Jaccard, H. Jindra, G. Weber, G. Wyss and J. Rabinowitz, Hel. Chem. Acta., 49, 807 (1966)



from NMR and chemical data that the product was 3-(N-phenylthiocarbamoyl)-2-phenyliminothiazolidine (3e).

As shown in Table I, the difference of pKa values for 2-amino-2-thiazoline (1a) and its secondary amine (1b) are larger than those of thiazole amines. The basicity of (1a) and (1b) are similar and a steric effect of N-methyl group in (1b) may be considered for the reaction. On the other hand, the aromatic secondary amine (1c) has a smaller pKa value than (1a, b) and is less basic, therefore, the nucleophillic activity of (1c) with (2a, b) is weaker than that of (1a, b) and a steric hindrance of phenyl group may be considered.

The reaction of (1c) with (2a) should provide either (3d) or its isomer (4d). Equimolar amounts (8.4 mmol) of (1c) and (2a) were refluxed for 2 hr. in dry toluene to give a yellow oily residue after evaporating the solvent which crystallized on cooling, m.p. 118-121° (crude). Recrystallization (EtOH) gave (3d), m.p. 123-124°. As shown in Table II, Rf value (TLC) of (3a) is larger than (4a) and (4c), and is close to the known (3a) and (3b).

As shown in Table III, NMR data of (3d) indicated a chemical shift of 4.82 ppm (t, J=7.0Hz, 2H) for N-CH₂, showing a lower field shift of 0.87 ppm than that of (1c). The authors^{1),6)} already pointed out that a low field shift of 0.80-0.90 ppm for $-N-CH_2$ at the position of 4 in the thiazolidine when an electron-withdrawing group such as thiocarbamoyl is attached at 3-N of 2-thiazoline. Therefore, (3d) seems to belong to the structure (3).

 ⁶⁾ Y. Yamamoto, R. Yoda and S. Kouda, Annual Report Kyoritsu Pharm. College, 18, 46(1973);
C. A., 81, 120524t (1974)

structure	pKa	solvent	temperature (°C)
$(1 a)$ $H \xrightarrow{N}_{H} S$	8.67 ± 0.03	H₂O	24
(1b) CH ₃ N H N S	8.66 ± 0.03	H ₂ O	24
$(1 c) \bigcirc H \xrightarrow{N} S$	7.20 ⁸⁾		
	2· 5 ⁷⁾	H₂O	20—25
	${\begin{array}{r} 5.40^{7} \\ 5.75 \ \pm \ 0.02 \end{array}}$	H₂O	26
H N S CH ₃	$\begin{array}{r} 5.81^{7)} \\ 5.75 \pm 0.02 \end{array}$	H₂O	24
CH ₃ N S CH ₃	$5.90~\pm~0.03$	H ₂ O	23

Table I pKa Values of (1a, b, c) and related compounds

7) M. Nagano, J. Tobitsuka, T. Matsui and K. Oyamada; Chem. Pharm. Bull. (Tokyo), 20, 2618 (1972)

8) H.Najer et al.; C. R. Acad. Sc. paris, t 260 (1965) 4343

When (3d) is refluxed in 15% H₂SO₄, it gave colorless crystals of m.p. 137° (EtOH). Its IR spectra showed a sharp strong band at 1700cm⁻¹ and was identified as 3–(N–methylthiocarbamoyl)thiazolidine–2–one $(5d)^{2}$ from IR and mixed m.p.. Therefore, the structure of (3d) was established to be 3–(N–methylthiocarbamoyl)–2–phenylimino–thiazolidine.

The reaction of 2-methylamino-2-thiazoline (1b) with phenylisothiocyanate (2c) may give 3-(N-phenylthiocarbamoyl)-2-methyliminothiazolidine (3c). An equimolar (1b) and (2c) reacted to give crystals (3) with m.p. 70-71° (EtOH). The Rf value

		Τί	able II Physical Consta	nts of Compounds			
	Compd.	m.p.	Found (Calcd.) (%)	IR (KBr disk)	10-21، میں 10-2).	TLC (Rf Value) cyclohexane 1	UHU
		(recrys.)	C : H : N	1650–1500cm ⁻¹	√max ^{mu} t¢ ∧ 10 /	CH _s COOC ₂ H ₅ i	C11.C13
	CH _S			1634			
3b		66°		1572	266. 0(180)		Ļ
	CH ₃	(EtOH)		1566	239. 0(sh)	0.03	0.40
	s			1605(s) 1568(s)			
3d		123—124°	52.74, 5.06, 16.96	1601(s) 1500(s)	275.0(183)		
	∩S∕N-©	(MeOH-H ₂ O)	(52.56, 5.21, 16.72)	1589(s)	246. 0(125)	0.65	0.36
	>						
	S = C			1620(s) 1550(s)	300.0(74)		
3c		7071°	52.51, 5.12, 16.69	1603(m) 1540(s)	270.0(119)	92 0	06 0
	CH ₃ CH ₃	(EtOH)	(52.56, 5.21, 16.72)	1590(m) 1530(s)	224.5(131)	2	60 •0
	U						
¢				1593(s) 1520(s)			
3e		116—117°	61.33, 4.61, 13.44	1570(s)	268.5(187)	0 63	07 0
	n 	(MeOH)	(61.31, 4.82, 13.41)	1540(s)	225.0(181)	8.0	0.40
	>						
4 c		150—152°		1615(w.sh)	331.5(76)		
	H H H	(CH ₃ CN)		1595(s . sh)	300.5(171)	0.36	0.10
	:			1580(s) 1570(s)			
				1516(s) 1510(s)			

No.23 (1978)

- 41 -

Table Ⅲ NMR Spectral Data by JNM-NH 100 (100MHz)

Con	npd.	N-CH ₂	CH ₂ -S	$=N-CH_8$	N-CH ₃	NH
3 b $\overset{CH_3}{\underset{H}{>}}N$	S -C N CH_2 CH_3 CH_3	4.65 t, J=7.0 Hz 2 H	3. 14 t, J=7.0 Hz 2 H	3.14 s, 3H		14.16 s, 1H
3 d CH ₃ N	S -C N CH_2 N S CH_2 O	4.82 t, J=7.0 Hz 2 H	3.10 t, J=7.0 Hz 2 H		3. 18 d, J=5. 0 Hz 3 H	11,98 s, 1H
3 c H	S N N CH ₂ CH ₂ CH ₂	4.82 t, J=7.0 Hz 2 H	3.14 t, J=7.0 Hz 2 H	3.14 s, 3H		14.16 s
3e H	S N-C-NCH ₂ N-S-CH ₂	4.85 t, J=7.0 Hz 2 H	3. 10 t, J=7.0 Hz 2 H			13.80
4 c H	S NCH₂ C _N LS-CH₂ H	3. 47 m	3. 23 m			9.22 10.32

The NMR Spectral data were measured at room temperature in $CDCl_3$. Chemical shifts are expressed in δ (ppm) with TMS as internal standard.

(TLC) is similar to (3d). NMR data showed a chemical shift of 4.82 ppm (t, J=7.0 Hz, 2H) for N-CH₂ which was same position with (3d) suggesting the structure (3). The hydrolysis of (3c) in 15% H₂SO₄ gave (1b) by elimination (2b). (2b) showed a characteristic $\nu_{N=C=S}$ absorption at 2130 cm⁻¹ (IR) and (1b) was identified by IR and as its picrate.

E. Cherbuliez *et al.*⁵⁾ heated (1c) and (2b) at 120°C for 2 hr. without solvent and the resulting crystals were recrystallized (MeOH), m.p. 112–113° and assigned as 2-(N-phenylthiocarbamoyl)phenylamino–2–thiazoline (4c).

The authors refluxed equimolar quantities of (1c) and (2b) in toluene the resulting product was recrystallized from EtOH to give colorless needles (3e), m.p. 116-117°. The product obtained by Cherbuliez's method gave an identical (3e) of m.p. 116-117° (EtOH). The Rf value (TLC) of the product (3e) is similar to those of (3c) and (3d) and the NMR data showed a chemical shift of 4.85 ppm (t, J=7.0Hz, 2H) which is

— 42 —

No.23 (1978)



Fig. I UV Absorption Spectra of (3b), (3c), (3d), (3e) and (4a) in propanol-2 for Spectroscopy (MERCK)

close to those of (3c) and (3d). The elemental analysis agreed with $C_{16}H_{15}N_3S_2$ and the IR spectra showed $\nu_{C=N}$ and $\nu \bigcirc$ absorptions at 1593 (s) and 1570 cm⁻¹ (s). The C=N absorption shifted to lower frequency due to the conjugation with phenyl group. As shown in Fig, I, UV absorption curves of (3c) and (3b) are similar. The 273 nm (ϵ =18300) band of (3d) corresponds with 270 nm (18300) of (3e). (3c), (3e) and (4a) all having N-phenylthiocarbamoyl group have an absorption at 300 nm. The absorption curve of (3e) is very different from that of (4a). As shown in Scheme I, (3e) was obtained from 1 mole of 1-amino-2-bromoethane (7) and 2 moles of (2b). As shown in Table IV, Mass spectral data indicated that the fragmentation of known (4c) and (3e) are quite different. (4c) showed M⁺ at m/e 237 (74.12%), (\bigcirc -N= C=S]⁺ at m/e 137 (base) and [C₆H₅]⁺ at m/e 77 (44.4%).

On the other hand, (3e) showed no M⁺ at m/e 313, but showed $[M^+- \bigcirc N=C=S]$ at m/e 177 (91.4%), $[\bigcirc -N=C=S]$ [†] at m/e 135 (base) and $[C_6H_5]^+$ at m/e 77 (base).

	Compd.	M+	M ⁺ -SH	$M^+-RN=C=S$		RN=C=S
3b	CH ₃ H N CH ₃ N C N S CH ₃	189 (84)	156 (1.1)	116 (base)	115 (88)	73 (18)
3 d		251 (70. 93)	_	178 (50)	177 (base)	73 (23. 58)
3c	⊖ N C N C N C N C N C N C N C N C N C N	251 (8.8)	_	116 (60. 3)	115 (base)	135 (61. 0)
3e		-	_	177 (91.4)	_	135 (base)
4 c	© _N -C _N LS	237 (74.1)	204 (8.1)	102 (33.8)	101 (10.6)	135 (base)

Table IV Mass Spectral Data m/e (Relative Intensity: %)

When (3e) was hydrolyzed with 15% H₂SO₄, it gave the products shown in Scheme II. As was the case for the hydrolysis of (3c) the liberation of (2b) took place quickly and (1c) was obtained as crystals when the aqueous layer was neutralized. The hydrolysis product aniline was detected by a color reaction. (2b) was identified with IR and (1c) was confirmed with IR and mixed m.p.

Conclusion

As for the reaction between (1) and (2), 2-amino-2-thiazoline (1a) as a primary amine (R=H) reacted with alkylisothiocyanate (2a) to give an unstable product (3a) which isomerized to (4a) upon heating. However, (1a) with arylisothiocyanate (2b) gave only stable (4c).

2-methylamino-2-thiazoline (1b) as a secondary amine $(R=CH_3)$ reacted with (2) (R'=alkyl or aryl) to give only (3) which did not isomerize upon heating.

2-phenylamino-2-thiazoline (1c) as an aromatic secondary amine $(R=C_6H_5)$ reacted with (2) (R'=alkyl or aryl) to give only (3d) or (3e).

No.23 (1978)



Scheme II

Experimental

Infrared spectra were obtained a Hitachi EPI-G3 spectrometer, ultraviolet spectra with a Shimadzu 200-S automatic recording spectrometer, nuclear magnetic resonance spectra with a Nihon Denshi TMS-OSG (70eV) and high mass resolution spectra with a Nihon Denshi JMS 01 SG-2 HI-resolution masspectrometer.

Typical experimental procedure is illustrated below;

3-(N-phenylthiocarbamoyl)-2-phenyliminothiazolidine (3e)

route A A toluene solution (30 ml) of 2-phenylamino-2-thiazoline (1 g, 0.0056 mol) and phenylisothiocyanate (1.52 g, 0.0112 mol) was refluxed for 2 hr. The toluene solu tion was evaporated to give crude (3e), which was recrystallized from ethanol to afford pure (3e).

route B To a mixture of cold 33% aq. KOH (100 ml) and benzene (100 ml) was added bromoethylamine HBr (33.2 g) with stirring. The benzene layer was separated and the aqueous layer was extracted with benzene (50 ml×2). To the combined benzene solution was added dropwise phenylthiocyanate (16 g) under ice-cooling, and then the mixture was refluxed for 2 hr. A viscous oily layer produced during the reaction was separated from the benzene layer and the benzene layer was extracted with H₂O. The aqueous layer was treated with conc. KOH (33%) to give white crystalline 2-ph-

enylamino-2-thiazoline (1c). The benzene layer evaporated *in vacuo* to leave a white solid (3e), which was recrystallized from methanol to afford pure (3e).

Hydrolysis of (3)

An aqueous HCl solution (15%, 20 ml) of (2a) was heated at about 110° . Just dissolving (3) in an aq. HCl solution, the oily substance was separated, and after being kept at room temperature for several hr. The solution was extracted with CHCl₃. The CHCl₃ layer was dryed over MgSO₄, and was evaporated to give oil (2b). The aqueous layer was treated with an aqueous NaHCO₃ to afford (1c, 0.25 g).