

Title	^2H nuclear magnetic resonance studies on biosynthesis : stereochemistry of the 5'-hydrogen atoms of griseofulvin derived from griseophenone B and 4-demethyldehydrogriseofulvin
Sub Title	
Author	佐藤, 良博(Sato, Yoshihiro) 小田, 泰子(Oda, Taiko) 斉藤, 肇(Saito, Hajime)
Publisher	共立薬科大学
Publication year	1978
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.23 (1978.) ,p.120- 121
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000023-0120

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.


^2H Nuclear Magnetic Resonance Studies on Biosynthesis: Stereochemistry of the 5'-Hydrogen Atoms of Griseofulvin derived from Griseophenone B and 4-Demethyldehydrogriseofulvin *

YOSHIHIRO SATO, TAIKO ODA and HAZIME SAITO (National Cancer Center Research Institute)

佐藤良博, 小田泰子, 齊藤 肇

We report the stereochemical fate of the deuterium atom at C-5' during the biosynthetic transformation of (1a) and (2a) by *Penicillium urticae* as studied by ^2H n.m.r. and mass spectrometry.

The two deuteriated tracer were synthesized as follows. Dehydrogenation of [5'- ^2H] griseofulvin with selenium oxide followed by demethylation as previously described



SCHEME. R^1 and R^2 correspond to the β - and α -configuration, respectively.

* J.C.S Chem. Comm., 1978, 135.

No.23 (1978)

gave 4-demethyl[5'-²H]dehydrogriseofulvin (2b). Treatment of (2b) with zinc-acetic acid afforded [5'-²H]griseophenone B (1b).

To a suspension in medium 2 (15 40-ml flasks, 15 ml per flask) of the mycelium obtained from 7-day-old shaken cultures of *penicillium urticae* in medium 1 (15 40-ml flasks, 20 ml per flask), was administered 15mg of [5'-²H]griseophenone B (1b). After a further 3 days griseofulvin (3b) was isolated from the broth. Similarly a tracer experiment with 4-demethyl[5'-²H]dehydrogriseofulvin (2b) afforded another deuteriated griseofulvin (3c) (Scheme). Since the ²H n.m.r. resonances of (3b) and (3c) are at the same position as that of the 5' α -signal of [5' α , 5' β -²H]griseofulvin (3d), the configuration of the deuterium atoms was unequivocally ascribed as 5' α .

Accordingly, the present ²H n.m.r. results confirm our previous conclusion that, in the feeding experiments involving [2-²H₃]acetate, the 5'-deuterium atom in biosynthetically deuteriated griseofulvin has the α -configuration.

The isotopic dilutions of (3b) and (3c), as calculated from mass spectroscopy, were 13 and 8, respectively, consistent with the postulated biosynthetic sequence that (2b) is at a closer stage than (1b) to the final product. Furthermore, it is interesting that the stereochemical course of the 5'-hydrogen atoms is the same as that in the microbial hydrogenation of dehydrogriseofulvin to griseofulvin (3a) by *Streptomyces cinereocrocatu*s NRRL 3443, suggesting the existence of similar enzyme systems in both microorganisms.