Title	The production of <sup>125</sup> Xe for medical use by the <sup>3</sup> He bombardment of natural tellurium
Sub Title	
Author	本間, 義夫(Honma, Yoshio)
	村上, 悠紀雄( Murakami, Yukio)
Publisher	共立薬科大学
Publication year	1978
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of
	Pharmacy). No.23 (1978. ) ,p.123- 123
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000023-0123

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって 保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

## The Production of <sup>125</sup>Xe for Medical Use by the <sup>3</sup>He Bombardment of Natural Tellurium \*

## YOSHIO HOMMA and YUKIO MURAKAMI

本間義夫,村上悠紀雄

<sup>133</sup>Xe (5.3 d,  $\beta^{-}$ ) has been extensively used to measure the ventilation of the lungs and blood flow to many organs and tissues. However, its physical properties are not optimal for in vivo studies. Recently, Hines et al. have compared <sup>133</sup>Xe and <sup>125</sup>Xe and have concluded that the half-life (17h), the decay mode (EC 100%), and the 188 keV and 284 keV  $\gamma$ -rays of <sup>125</sup>Xe make it a particularly useful diagnostic scanning agent, giving a much lower radiation dose rate for a given count rate than the more readily available <sup>133</sup>Xe. The production method proposed by Hines et al. produces <sup>125</sup>Xe in a yield of 11 mCi/ $\mu$ Ah. However, for satisfactory yields this method requires bombardment by 31 MeV protons, which are not attainable with the compact medical cyclotrons suitable for routine production. In this work, in order to find out the optimum bombardment conditions for  $\alpha$  and <sup>3</sup>He reactions producing <sup>125</sup>Xe, the authors have measured the excitation functions and thick-target yield curves from natural tellurium. The Te (<sup>3</sup>He, xn) <sup>125</sup>Xe and the Te ( $\alpha$ , xn) <sup>125</sup>Xe reactions have cross section peaks of 117 and 51 mb at 35 and 40 MeV, respectively. The thick-target yields of 125Xe at 40 MeV <sup>3</sup>He and  $\alpha$  were 280 and  $40\mu$ Ci/ $\mu$ Ah, respectively. That is to say, for the production of <sup>125</sup>Xe, the <sup>3</sup>He bombardment is more advantageous than  $\alpha$  bombardment. The bombardment of natural ellurium with 40 MeV <sup>3</sup>He particles also produced <sup>123</sup>Xe, <sup>123</sup>I and <sup>130</sup>I in yields of 220, 85 and  $213\mu$ Ci/ $\mu$ Ah, respectively. When <sup>125</sup>Xe was produced for medical application, therefore, a 21-hr cooling period was required before the <sup>125</sup>Xe separation to allow the <sup>123</sup>Xe to decay. The advantages of the proposed method are that usable quantities of <sup>125</sup>Xe can be produced with a relatively low bombardment energy and an inexpensive natural tellurium target.

<sup>\*</sup>本総説は Int. J. appl. Radiat. Isotopes. 28, 738 (1977) に発表。