Title	Studies on monoamine oxidase (IV) : monoamine oxidase activity in rat brain and estrous cycle
Sub Title	
Author	木村, 都(Kimura, Miyako)
	中村, 悦郎(Nakamura, Etsuro)
Publisher	共立薬科大学
Publication year	1978
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of
	Pharmacy). No.23 (1978.) ,p.132- 132
JaLC DOI	
Abstract	
Notes	学会講演要旨
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000023-0135

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって 保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

No.23 (1978)

は1,18hrで増加し、cGMPは2hrまでは著明に減じ、18hrでも回復しない。従ってcAMP /cGMPは2~18hrで増加する。寒冷ストレス負荷による脳内及び副腎髄質のTH活性の変動 はcAMPの影響下にあると考えられる。TH活性の殆んど認められない副腎皮質のcAMP, cGMPの前述の変化については今後ステロイド分泌機構との関連を追求したい。

Studies on Monoamine Oxidase (IV): Monoamine Oxidase Activity in Rat Brain and Estrous Cycle

MIYAKO KIMURA and ETSURO NAKAMURA

木村 都,中村悦郎

〔第51回日本薬理学会総会 仙台(1978年3月)で発表〕

The activity of monoamine oxidase (MAO) during different phases of the estrous cycle was studied using seven brain regions from S.D. strain rats. The rats were maintained on a 12 hr light-dark schedule. MAO activity of homogenate of each tissue was assayed by isotope dillution method using ¹⁴C-tyramine as substrate. In most regions of the brain, MAO activity was highest in proestrus (P), subsequently fell in estrus I (EI), rose again in estrus II (EII), and turned to a lower level during diestrus. Especially in the hypothalamus, MAO activity in P showed highly a significant difference (p < 0.01) from that in diestrus. MAO activity in the hypothalamus of rats 2 weeks after ovariectomy was lower than that in P phase of nonoperated rats. When estradiol (0. 2mg/kg/day for 3 days) was given s.c. to ovariectomized rats, MAO activity in the hypothalamus was depressed significantly, whereas progesterone (4mg/kg /day for 3 days) showed tendency toward restration of the activity depressed to the level in P. Estradiol together with progesterone also inhibited the activity to the same extent as the former alone. Plasma levels of estrogens were considered to be highest in P. The peak of the activity appeared almost simultaneously with the so-called critical period. Thus, no direct relationship between the fluctuation of MAO activity in the hypothalamus and different levels of endogeous steroids is recognized.

Fluctuation in the Levels of Cyclic AMP and Cyclic GMP in Rat Brain during Estrous Cycle

JUNKO ONO, MIYAKO KIMURA and ETSURO NAKAMURA

小野純子,木村 都,中村悦郎

〔第51回日本薬理学会総会 仙台(1978年3月)で発表〕

In connection with the effects of sexual steroids, fluctuations in the levels of the