

Title	Viscometric analysis of effects of cytochalasans on in vitro polymerization and depolymerization of microtubules
Sub Title	
Author	佐藤, 良博(Sato, Yoshihiro) 斎藤, 祐子(Saito, Yuko) 手塚, 貴子(Tezuka, Takako) 関田, 節子(Sekita, Setsuko) 義平, 邦利(Yoshihira, Kunitoshi) 名取, 信策(Natori, Shinsaku)
Publisher	共立薬科大学
Publication year	1982
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.27 (1982.) ,p.46- 48
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000027-0046

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

VISCOMERTIC ANALYSIS OF EFFECTS OF CYTOCHALASANS ON IN VITRO POLYMERIZATION AND DEPOLY- MERIZATION OF MICROTUBULES*

YOSHIHIRO SATO, YUKO SAITO, TAKAKO TEZUKA, SETSUKO SEKITA,
KUNITOSHI YOSHIHARA and SHINSAKU NATORI

佐藤良博, 斎藤祐子, 手塚貴子, 関田節子**
義平邦利,** 名取信策**

This paper describes studies which clarify that 19 cytochalasins, including 6 natural cytochalasins, 2 natural aspochalasins, 7 natural chaetoglobosins and 4 their acetates, except for CA do not show any marked inhibitory effects on the association of porcine brain tubulin and also on the dissociation of the microtubule, analyzed by viscosity measurement.

Results of the cytochalasins on the inhibitions of *in vitro* polymerization and

Table I. The Effects of Cytochalasins for the *in Vitro* Microtubule Polymerization and the *in Vitro* Depolymerization of Microtubule determined by Viscometry

Substance	$(a-b)/a$	$(a'-b')/a'$
Cytochalasin A (CA)	0.37 ****a)	0.35 ****a)
Cytochalasin B (CB)	0.05 *	0.00 *
Cytochalasin C (CC)	0.06 *	0.10 *
Cytochalasin D (CD)	-0.03 *	0.00 *
Cytochalasin E (CE)	0.01 *	0.03 *
Cytochalasin H (CH)	-0.08 *	0.05 *
Aspochalasin B (AsB)	0.06 *	0.11 **
Aspochalasin D (AsD)	0.13 **	0.12 **
Chaetoglobosin A (ChA)	0.23 ***	0.16 **
Chaetoglobosin A-Ac (ChA-Ac)	0.19 **	0.06 *
Chaetoglobosin B (ChB)	0.15 **	0.07 *
Chaetoglobosin C (ChC)	-0.03 *	0.01 *
Chaetoglobosin D (ChD)	0.15 **	0.10 **
Chaetoglobosin E (ChE)	-0.05 *	-0.03 *
Chaetoglobosin E-Ac (ChE-Ac)	0.02 *	0.09 *
Chaetoglobosin F (ChF)	0.01 *	0.03 *
Chaetoglobosin F-Ac (ChF-Ac)	0.05 *	0.08 *
Chaetoglobosin J (ChJ)	0.09 *	0.10 **
Chaetoglobosin J-Ac (ChJ-Ac)	0.10 **	0.10 **

a) The values of $(a-b)/a$ and $(a'-b')/a'$ were ranked by the number of the asterisk(*) : *, 0.10 > $(a-b)/a$ and $(a'-b')/a'$, **; 0.20 > $(a-b)/a$ and $(a'-b')/a' \geq 0.10$, ***; 0.30 > $(a-b)/a$ and $(a'-b')/a' \geq 0.20$, and ****; 0.40 > $(a-b)/a$ and $(a'-b')/a' \geq 0.30$.

* 本報は *J. Pharm. Dyn.*, 5, 418 (1982) に報告.

** 国立衛生試験所

depolymerization of microtubules were shown in Table I. The structures of cytochalasins are shown in Fig. 1. The dose-response curves of the inhibition of the polymerization of microtubule proteins by CA, CC, and ChA were also shown in Fig. 2.

Firstly, the results given in Table I show that no cytochalasins except for CA have activities in the inhibition of microtubule polymerization *in vitro* and of the depolymeriza-




Fig. 1. Structures of Cytochalasins
a) Cytochalasasin A and B, b) Chaetoglobosin A and D,
c) Aspochalasin B and D.




Fig. 2. Dose-response Curves of the Inhibition of Polymerization of Microtubule Proteins by Vinblastine (Δ), Colchicine (\blacktriangle), Griseofulvin (\circ), Daunomycin (\square), Cytochalasin A (\blacksquare), and C (\odot), and Chaetoglobosin A. Each point is the mean of at least two experiments.

tion of polymerized tubulin (microtubules) *in vitro*. The values of $(a-b)/a$ and $(a'-b')/a'$ of CA were 0.37 and 0.35, respectively, and those of ChA follow by 0.23 and 0.16, respectively. All other cytochalasans marked by the asterisk of * or ** have the values of $(a-b)/a$ and $(a'-b')/a'$ less than 0.10 or 0.20, respectively, indicating negligible inhibitory activities on the polymerization of porcine microtubules *in vitro*. Fig. 2 represents the dose-response curves of three cytochalasans : CA, CC, and ChA with those of vinblastine, colchicine and griseofulvin which are known as the sensitive antimitotic drugs, and daunomycin which is also reported as an active component for representative antimitotic drugs, and daunomycin which is also reported as an active component for perturbing the polymerization of tubulin and microtubule assembly. The results obtained in this experiment indicate that, under the conditions in which vinblastine and colchicine inhibit the polymerization of tubulin into microtubules in low concentration, CA showed a slight but distinct effect. However, its activity was lower than that of griseofulvin and higher than those of daunomycin and ChA. The dose-response curve of CC represents the activities of all other cytochalasans which showed small $(a-b)/a$ values. On the other hand, the micromolar concentrations of drugs which show the value of 0.2 in reference to $(a-b)/a$ were calculated as follows : vinblastine, 0.04 ; colchicine 0.33 ; griseofulvin, 23 ; cytochalasin A, 44 ; daunomycin, 110 ; chaetoglobosin A, 190.

Although data of the present studies were evaluated only on the basis of the results of the viscometric analyses, the results indicate that the *in vitro* target protein of the cytochalasans is not the tubulins or microtubules. In conclusion, our results could be evaluated in connection with the studies of Yahara et al. which suggested the interaction of cytochalasans with actin.