Title	Spectrophotometric determination of palladium with a new chelating reagent, N, N-dimethyl-N'-(4-methyl-5-nitro-2-thiazolyl) thiourea
Sub Title	
Author	与田, 玲子(Yoda, Reiko)
Publisher	共立薬科大学
Publication year	1982
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.27 (1982.),p.57- 57
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000027-0057

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

**Spectrophotometric Determination of Palladium with a
New Chelating Reagent, N,N-Dimethyl-N'-(4-methyl-5-nitro-2-thiazolyl) thiourea***

REIKO YODA

与田玲子

A new spectrophotometric method for the determination of palladium with *N,N*-dimethyl-*N'*-(4-methyl-5-nitro-2-thiazolyl)thiourea (DMNT) is proposed. DMNT instantly forms a yellow chelate with Pd (II) at pH 2 at room temperature, which is readily extracted into organic solvents such as chloroform and shows a higher absorption maximum ($\epsilon=40,400$) at 413 nm. The ratio of Pd(II) to reagent in the chelate is 1 : 2 as determined by the widely used method and also verified by synthesis of the authentic Pd-chelate. The chelate conforms well with the Lambert-Beer's law over a wide concentration range (3.3—22 μg in 10 ml chloroform). The optimum concentration range of palladium for spectrophotometry by the Ringbom plot is 5.6—18.6 μg in 10 ml of chloroform solution. This method gives a good reproducibility, high sensitivity and high accuracy in the presence of many foreign ions.

* 本報告は, Mikrochimica Acta [Wien] 1982 II 271-278 に発表.