

| | |
|------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Title | Substituent effects of potassium phenoxides on the carboxylation of indene by carbon dioxide |
| Sub Title | |
| Author | 森, 久和(Mori, Hisakazu) 峯田, 一幸(Mineta, Kazuyuki) 管, 孝男(Kan, Takao) |
| Publisher | 共立薬科大学 |
| Publication year | 1983 |
| Jtitle | 共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.28 (1983.) ,p.94- 94 |
| JaLC DOI | |
| Abstract | |
| Notes | 抄録 |
| Genre | Technical Report |
| URL | https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000028-0094 |

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Substituent Effects of Potassium Phenoxides on the Carboxylation of Indene by Carbon Dioxide*

Hisakazu MORI, Kazuyuki MINETA,** and Takao KWAN***

森 久和, 峯田一幸,** 菅 孝男***

The carboxylation of indene by carbon dioxide was investigated in the presence of substituted phenoxide in DMF. The reaction took place rapidly at 0°C and was apparently completed within about ten minutes. Indene-3-carboxylic acid was found to be formed in 42—98% yields from indene, depending on the substituents (*p*-OC₄H₉, *p*-OCH₃, *p*-CH₃, H, *p*-Cl, *m*-Cl, *p*-CN, and *m*-NO₂) of the phenoxides. The substituent effect upon the interaction of carbon dioxide with substituted phenoxides was observed. And it was found that the yield of indene-3-carboxylic acid increased with the substituents of negative σ values, giving rise to a linear relation between the logarithm of the equilibrium constants of carboxylation of indene and the σ values. Relative rates of reaction were compared with the various substituted phenoxides mentioned above. The mechanism of reaction is briefly discussed.

* 本報告は *Chem. Pharm. Bull.*, 31, 3002—3008 (1983) に発表

** 三菱化成

*** 帝京大学薬学部