Title	Mechanism of stigmasterol dealkylation in insect
Sub Title	
Author	藤本, 善徳(Fujimoto, Yoshinori) 木村, 美記(Kimura, Miki) 鷹巣, 彰弘(Takasu, Akihiro) Khalifa, Fathy A. M.(Morisaki, Masuo) 森崎, 益雄(Ikekawa, Nobuo) 池川, 信夫
Publisher	共立薬科大学
Publication year	1984
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.29 (1984.),p.36- 36
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000029-0036

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Mechanism of Stigmasterol Dealkylation in Insect*

Yoshinori FUJIMOTO**, Miki KIMURA**, Akihiro TAKASU**,
Fathy A. M. KHALIFA**, Masuo MORISAKI and Nobuo IKEKAWA**

藤本善徳**, 木村美記**, 鷹巣彰弘**, ファシー・カリファ***,
森崎益雄, 池川信夫**

Deuterated stigmasterols were chemically synthesized and fed to silkworm larvae. GC-MS analysis of metabolites, cholesterol and desmosterol indicates the metabolic transformations shown below, involving the migration of 25-hydrogen to C-24 position during stigmasterol dealkylation.

The 22,24(28)-dienes were shown to be converted to the 22,24-diene, desmosterol and cholesterol. All these results strongly support the mechanism depicted in the following scheme.

* 本報告は *Tetrahedron Lett.*, 25, 1501 (1984) に発表

** 東京工業大学理学部

*** エジプト国, 国立カイロ大学