

Title	Products of the nitration of 2-thiazolylureas and 2-thiazolylthioureas
Sub Title	
Author	与田, 玲子(Yoda, Reiko) 山本, 有一(Yamamoto, Yuichi) 岡田, 智子(Okada, Tomoko) 松島, 美一(Matsushima, Yoshikazu) 飯高, 洋一(Iitaka, Yoichi)
Publisher	共立薬科大学
Publication year	1984
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.29 (1984.) ,p.45- 46
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000029-0045

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Products of the Nitration of 2-Thiazolylureas and 2-Thiazolylthioureas*

Reiko YODA, Yuichi YAMAMOTO, Tomoko OKADA,
Yoshikazu MATSUSHIMA and Yoichi IITAKA**

与田玲子, (故)山本有一, 岡田智子,
松島美一, 飯高洋一**

Our recent studies have shown that 2-thiazolylthioureas are a fruitful source of chelating agents which can be used as reagents for spectrophotometric determination of metal ions. Introduction of a nitro group in the 5-position of the thiazole moiety of 1,1-dimethyl-3-(4-methyl-2-thiazolyl)thiourea caused bathochromic and hyperchromic effects on the absorption bands of the thioureas and their metal chelates. This finding prompted us to prepare the corresponding nitro compounds of the other series of 2-thiazolylureas and 2-thiazolylthioureas.

Nitration was performed by a dropwise addition of fuming nitric acid to a concentrated




Chart 1




Chart 2

* 本報告は *Chem. Pharm. Bull.*, 32, 3483—3492 (1984) に発表

** 東京大学薬学部

sulfuric acid solution of a urea or a thiourea at low temperatures. Nitration products of 1-methyl-3-(4-methyl-2-thiazolyl)-urea (1) and -thiourea (2) were their 5-nitrothiazolyl derivatives (3 and 4, respectively). With an excess of nitric acid, 1-methyl-3-(4-methyl-5-nitro-2-thiazolyl)-1-nitrourea (5) was obtained from 1, whereas corresponding 1-nitrosourea (6) from 2. From the nitration of 3-(4-methyl-2-thiazolyl)thiourea (7), a pale yellow compound (9) was obtained with the 5-nitrothiazolyl derivative of 7 (8). Unlike other nitrated thioureas, 9 did not form a colored Cu(II) chelate and was stable to acid, alkali, and heat. It was concluded to be 6-methyl-2-nitroiminothiazolo[3,2-b][1,2,4]-thiadiazole from studies of its physicochemical properties, chemical reactions, and crystallography. Corresponding compounds were obtained from other N-(4-alkyl-2-thiazolyl)-thioureas. Chemical reactions studied are summerized in Chart 1 and 2.