

Title	Effects of acute and chronic treatments with atenolol and propranolol on cardiovascular responses to handling stress in spontaneously hypertensive rats
Sub Title	
Author	工藤, 善隆(Kudo, Yoshitaka) 曾我部, 博文(Sokabe, Hirofumi) 川島, 紘一郎(Kawashima, Koichiro)
Publisher	共立薬科大学
Publication year	1984
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.29 (1984.) ,p.54- 54
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000029-0054

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

**Effects of Acute and Chronic Treatments with Atenolol and
Propranolol on Cardiovascular Responses to Handling Stress
in Spontaneously Hypertensive Rats***

Yoshitaka KUDO**, Hirofumi SOKABE** and Koichiro KAWASHIMA

工藤善隆**, 曾我部博文**, 川島紘一郎

Although the mechanism of antihypertensive action of β -adrenergic blocking drugs (β -blockers) is not known, a theoretical advantage of cardioselective β -blockers over nonselective ones has been proposed in the treatment of hypertension. To study this hypothesis, we examined cardiovascular responses to handling stress in spontaneously hypertensive (SHR) rats after a single (100 mg/kg) and multiple oral treatments (100 mg/kg per day for 17 d) with either atenolol or propranolol. Atenolol and propranolol markedly suppressed the tachycardia induced by handling stress after acute and chronic administration. Resting mean arterial pressure (MAP) was reduced by acute and chronic atenolol treatment, but not by propranolol. Stress-induced increase in MAP was significantly reduced by chronic treatment with propranolol, whereas no consistent effects were observed with atenolol. Acute treatment with guanethidine (30 mg/kg) markedly reduced the rise in MAP induced by stress. These results suggest that suppression of cardiac function by β -blockers does not always attenuate the rise in MAP induced by stress, thus cardioselective β -blockers might not confer any further reduction of the blood pressure increase due to sympatho-adrenal excitation. Inhibition of stress-induced MAP rise by propranolol could be mediated by a modulation of the catecholamine release.

* 本報告は *J. Pharm. Dyn.*, 6, 729—736 (1983) に発表

** 自治医大・薬理