

Title	The structure of panaxan B, a hypoglycemic glycan of panax ginseng
Sub Title	
Author	友田, 正司(Tomoda, Masashi) 嶋田, 和代(Shimada, Kazuyo) 今野, 長八(Konno, Chohachi) ヒキノ, ヒロシ(Hikino, Hiroshi)
Publisher	共立薬科大学
Publication year	1985
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.30 (1985.) ,p.64- 65
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000030-0064

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

The Structure of Panaxan B, A Hypoglycemic Glycan of *Panax ginseng**

Masashi TOMODA, Kazuyo SHIMADA, Chohachi KONNO** and Hiroshi HIKINO**

友田正司, 嶋田和代, 今野長八**, ヒキノヒロシ**

The roots of *Panax ginseng* C.A. MEYER has been utilized as an elixir in Oriental medicine. This crude drug has long been known to possess the ability to lower blood sugar level in normal and experimentally-induced hyperglycemic animals. Recently, we have identified the hypoglycemic peptidoglycans, panaxan A, B, C, D and E, in ginseng and we have determined the partial structure of one of the main peptidoglycans, panaxan A. The present paper deals with the structural study of panaxan B, another main peptidoglucan in ginseng.

When panaxan B was subjected to gel chromatography on Sephacryl S-500, a value of *ca* 1 800 000 for the molecular weight was obtained. Quantitative determination showed that the component sugar of panaxan B was D-glucose (95.9%) and a small amount of peptide moiety was present (0.7%).

Panaxan B exhibited a high positive specific rotation ($[\alpha]_D + 196^\circ$) and its ¹H-NMR spectrum disclosed two anomeric hydrogen signals at δ 4.88 (doublet, *J* 3 Hz) and 5.24 (doublet, *J* 3 Hz), and the ratio of their integrals was about 3 : 1. Thus α -linkage of all D-glucose units was deduced.

Among nine signals at δ 60.6, 65.3, 69.6, 70.2, 71.7, 73.3, 81.5, 97.9 and 99.6 observed in the ¹³C-NMR spectrum of panaxan B, it was evident that those at δ 60.6 and 65.3 were attributable to the C-6 carbons, those at δ 73.3 and 81.5 ascribable to the C-3 carbons, and those at δ 97.9 and 99.6 assignable to the C-1 carbons. Therefore, it was concluded that α -glucose units are linked at the 1, 3 and 6 positions in panaxan B.

Methylation of panaxan B by the Hakomori's method afforded the fully methylated derivative which was hydrolysed with dilute H₂SO₄ in HOAc. The products were converted into alditol acetates and analysed by GC-MS. As a result, 2,3,4,6-tetra-*O*-methyl-D-glucose, 2,3,4-tri-*O*-methyl-D-glucose and 2,4-di-*O*-methyl-D-glucose were identified and their approximate molar ratio was 1.0 : 1.8 : 1.0.

Periodate oxidation of panaxan B consumed 1.47 mol of periodate per mol of component anhydro sugar unit with liberation of 0.83 mol of formic acid. When the periodate-oxidized product was reduced, hydrolysed and analysed, the residual glucose was obtained in a yield of 31.0%.

Accumulated data demonstrated that panaxan B is mainly composed of α -1→6

* 本報告は *Phytochemistry*, 24, 2431—2433 (1985) に発表

** 東北大学薬学部

linked D-glucopyranose units and possesses branches linked in part at the C-3 position. The ratio of terminals, branching positions and intermediate units should be *ca* 1 : 1 : 1.8 on the average. Judging from the above facts, there are more branchings in panaxan B than panaxan A.