Title	Pyrolytic products of N-alkyl-N'-(4-methyl-2-thiazolyl)-S-methylisothioureas
Sub Title	
Author	与田, 玲子(Yoda, Reiko) 山本, 有一(Yamamoto, Yuichi) 岡田, 智子(Okada, Tomoko) 松島, 美一(Matsushima, Yoshikazu)
Publisher	共立薬科大学
Publication year	1985
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.30 (1985.),p.91- 91
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000030-0091

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

**Pyrolytic Products of N-Alkyl-N'-(4-methyl-2-thiazolyl)-
S-methylisothiourreas***

Reiko YODA, Yuichi YAMAMOTO, Tomoko OKADA and Yoshikazu MATSUSHIMA

与田玲子, 山本有一, 岡田智子, 松島美一

We reported previously that pyrolysis of N,N-dimethyl-N'-(4-methyl-2-thiazolyl)-S-methylisothiourrea at 155°C for 14 h gave N,N-dimethyl-N'-(3,4-dimethyl-2(3H)thiazolylidene)thiourea as the main product. The pyrolysis proceeded smoothly without any solvent. In the pyrolysis of other N,N-dialkyl-N'-(2-thiazolyl)-S-methylisothiourreas, a similar migration of the methyl group from the isothiourrea sulfur atom to the thiazole nitrogen atom was observed.

In contrast, the pyrolysis of an N-nonomethyl analog, N-methyl-N'-(4-methyl-2-thiazolyl)-S-methylisothiourrea (**1a**), under the same conditions gave N-methyl-N',N''-bis-(4-methyl-2-thiazolyl)guanidine (**2a**) as a product. The S→N methyl migrated compounds such as N-methyl-N'-(3,4-dimethyl-2(3H)thiazolylidene)thiourea and N-methyl-N'-(3,4-dimethyl-2-thiazolyl)-N''-(4-methyl-2-thiazolyl)guanidine (**6a**) were not formed in the pyrolysis. Similarly, a pyrolytic product of an N-ethyl analog of **1a**, N-ethyl-N'-(4-methyl-2-thiazolyl)-S-methylisothiourrea (**2b**), was identified as an N-ethyl analog of **2a**, N-ethyl-N',N''-bis(4-methyl-2-thiazolyl)guanidine (**2b**).

The difference in the pyrolytic behavior between the N,N-dialkyl- and N-monoalkyl-N'-(2-thiazolyl)-S-methylisothiourreas were interpreted in terms of the basicity of the thiazole nitrogen atom and the predominant chemical species.

* 本報告は *Heterocycles*, **23**, 2339 (1985) に発表