

Title	Effects of dexamethasone on metallothionein induction by Zn, Cu and Cd in Chang liver cells
Sub Title	
Author	小林, 静子(Kobayashi, Shizuko) 岡田, 貴子(Okada, Takako) 木村, 正己(Kimura, Masami)
Publisher	共立薬科大学
Publication year	1986
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.31 (1986.) ,p.59- 60
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000031-0059

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Effects of Dexamethasone on Metallothionein Induction by Zn, Cu and Cd in Chang Liver Cells*

Shizuko KOBAYASHI, Takako OKADA and Masami KIMURA**

小林静子, 岡田貴子, 木村正己**

Karin et al. have described that human MT are encoded by a multi-gene family containing at least 14 closely related genes and pseudogenes. Two of these genes, MT-I_A and MT-II_A, encode functional protein which show each other different induction response to heavy metal and glucocorticoids. The differential response is due to functional differences of the respective promoter/regulatory regions of the genes; the MT-I_A promoter response only to Cd²⁺, but MT-II_A promoter is responsive to Cd²⁺, Zn²⁺ and glucocorticoids. Moreover, they have defined two separate regulatory DNA elements for the MT-II_A gene induction which mediate the induction of the gene by heavy metals and by glucocorticoid hormones. The element responsible for induction by glucocorticoid hormones is coincident with the DNA-binding site for the glucocorticoid hormone receptor. These results clearly indicate that two inducers, heavy metals and glucocorticoid hormones, promote MT gene expression by mechanisms which are independent of one another.

We are interested in the extent to which particular inducers affect the induction of MT. In this report we demonstrate the effects of dexamethasone on maximum MT induction by Zn²⁺, Cu²⁺ and Cd²⁺ in Chang liver cells. Our data show that dexamethasone, has an additive effect on the maximum MT accumulation induced by Zn²⁺, Cu²⁺ or Cd²⁺, supporting the interpretation that expressions of mechanism of MT gene expression by heavy metals and dexamethasone work independently of one another.

Metallothioneins (MTs) were induced in Chang liver cells by the metals, Zn, Cu and Cd, and the glucocorticoid hormone, dexamethasone. When 116 μ M Zn, 32 μ M Cu and 18 μ M Cd, and 10⁻⁷M dexamethasone, respectively, were administered for 9 h, MTs induced by each inducer in the cells reached maximum levels. The maximum accumulation of MT level induced by dexamethasone was the lowest of the four inducers investigated; the levels induced by Zn, Cu and Cd were 4.7, 1.2 and 1.5 times of that induced by dexamethasone.

When dexamethasone was added to the cells together with the heavy metals (Zn, Cu and Cd), dexamethasone had an additive effect on the maximum MT accumulations induced by heavy metals as compared to when induction was conducted using one of

* 本報告は *Chem-Biol. Interactions*, 55, 347—356 (1985) に発表.

** 産医研.

heavy metals alone or by dexamethasone alone. However, dexamethasone did almost not effect the metal accumulations in the cells, although the maximum MT levels induced by heavy metal increased by dexamethasone. These results suggest that the process of MT induction by heavy metals and that by dexamethasone are independent of one another.

When dexamethasone was added to the cells together with a high concentration of Cu ($32\ \mu\text{M}$) induced the maximum MT accumulation, Cu transport into the cells decreased by 20—40% of that into non-treated cells, which was statistically significant.

In this experiment, our data suggest that MT genes possess a differential gene expression mechanism by functional differences of MT : detoxification of heavy metals and regulation of essential heavy metals.