Title	Enkephalin but not morphine modulates the motor activity in the frog spinal cord in vitro
Sub Title	
Author	鈴木, 岳之(Suzuki, Takeshi) 岡, 淳一郎(Oka, Junichiro) 永野, 伸郎(Nagano, Nobuo) 福田, 英臣(Fukuda, Hideomi)
Publisher	共立薬科大学
Publication year	1986
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.31 (1986.),p.74- 74
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000031-0074

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Enkephalin But Not Morphine Modulates The Motor Activity in the Frog Spinal Cord *In Vitro*

Takeshi SUZUKI, Jun-ichiro OKA*, Nobuo NAGANO* and Hideomi FUKUDA*

鈴木岳之, 岡 淳一郎,* 永野伸郎,* 福田英臣*

1. In the isolated frog spinal cord, methionine-enkephalin (ME, up to $3 \times 10^{-5}M$) hyperpolarized the resting ventral and dorsal root potentials.
2. These hyperpolarizations remained even under Ca^{2+} -free conditions.
3. ME depressed the fast component of the electrically stimulated spinal reflex and enlarged the following depolarizing component.
4. Morphine ($10^{-4}M$) had no apparent ME-like effects.
5. ME may directly and indirectly affect the motoneuron and modulate the spinal motor activity in the frog spinal cord.

本報告は *Comp. Biochem. Physiol.*, Vol. 83 C (2), 245—251 (1986) に発表.

* 東京大学薬学部毒性薬理学教室.