

Title	Porphyrins with metal-chelating groups in the peripheral region. II. meso-tetra [2-(8-hydroxyquinoly)] porphine
Sub Title	
Author	菅田, 節朗(Sugata, Setsuro) 松島, 美一(Matsushima, Yoshikazu)
Publisher	共立薬科大学
Publication year	1987
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.32 (1987. ) ,p.44- 44
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	<a href="https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000032-0044">https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000032-0044</a>

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

**Porphyrins with Metal-chelating Groups in the Peripheral Region. II.**  
***meso*-Tetra[2-(8-hydroxyquinolyl)]porphine\***

Setsuro SUGATA and Yoshikazu MATSUSHIMA

菅田節朗, 松島美一

It is well known that many natural and synthetic porphyrins accumulate *in vivo* in neoplastic tissues, though the affinity for the tissue seems to be lost in metalloporphyrins. Porphyrins with metal-chelating groups in the peripheral region can be obtained as the free base form of the porphyrin labeled with radioactive or nonradioactive metal ions at the periphery. The potential uses of such porphyrins in medical sciences and analytical chemistry are many.

Previously we reported the synthesis of *meso*-tetra[5-(8-hydroxyquinolyl)]porphine as a model of such porphyrins. We have now prepared one of its isomers, *meso*-tetra[2-(8-hydroxyquinolyl)]porphine by the Rothmund method. Infrared and mass spectra confirmed the structure.

The compound was soluble in dimethylformamide (DMF) but practically insoluble in water. Addition of Cu(II) perchlorate to the DMF solution resulted in biphasic spectral changes, an instantaneous change and a subsequent slow change. The spectrum after the first step showed the formation of the free base porphyrin with metal-chelated 8-quinolinol moieties. The second step was observable in the presence of a large excess of Cu(II) and was ascribed to the formation of the Cu(II)-porphyrin with Cu(II)-chelated 8-quinolinol. Addition of perchloric acid to the solution of the metalloporphyrin gave rise to the spectrum assignable to the Cu(II) porphyrin with unchelated 8-quinolinol.

---

\* 本報告は *Chem. Pharm. Bull.*, 35, 2623 (1987) に発表.