Title	Chemical and mutagenic properties of α-phosphonooxynitrosamines
Sub Title	
Author	望月, 正隆(Mochizuki, Masataka) 五十嵐, 敦子(Igarashi, Atsuko) 鈴木, 厚子(Suzuki, Atsuko) 関口, 奈保子(Sekiguchi, Naoko) 安生, 孝子(Anjo, Takako) 岡田, 正志(Okada, Masashi)
Publisher	共立薬科大学
Publication year	1987
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.32 (1987.) ,p.48- 48
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000032-0048

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって 保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Chemical and Mutagenic Properties of *a*-Phosphonooxynitrosamines*

Masataka Mochizuki, Atsuko Ikarashi, Atsuko Suzuki, Naoko Sekiguchi, Takako Anjo** and Masashi Okada**

望月正隆,五十嵐敦子,鈴木厚子,関口奈保子,安生孝子**,岡田正志**

The chemical and mutagenic properties of the products of solvolysis of α -acetoxynitrosamines in phosphate buffer were investigated. α -Acetoxynitrosamines decomposed in two ways: O-acyl fission yielded α -hydroxynitrosamines, which decomposed into aldehydes and alcohols, while O-alkyl fission gave a resonance hybrid of α -N-nitrosocarbonium and -iminium ions, which was trapped with phosphate and afforded α -phosphono-Formation of α -phosphonooxynitrosamines was dependent on the strucoxynitrosamine. ture of α -acetoxynitrosamines; those with a secondary α -phosphonooxy group, including cyclic nitrosamines, were easily formed, while among those with a primary phosphonooxymethyl group, only those with an alkyl group containing a branched α -carbon as isopropyl, sec-butyl and tert-butyl were isolated. They were good substrates of alkaline phosphatase and showed a nuclear magnetic resonance spectrum due to the presence of a phosphorus atom. They were decomposed by acid catalysis, and the rate was dependent on the structure. They were directly mutagenic in bacterial tester strains, except for a compound with a *tert*-butyl group. The activity was similar or stronger in Salmonella typhimurium TA 1535 and much weaker in Escherichia coli WP 2 and WP 2 hcr⁻ than those of α -acetoxynitrosamines. Stability in neutral aqueous solution and the strong mutagenicity of α -phosphonooxynitrosamines suggested their possible involvement in metabolic activation as a precursor of α -hydroxynitrosamines, and also in the organotropic carcinogenicity of N-nitrosodialkylamines as a transport form.

^{*} 本報告は IARC Scientific Publications, No. 84, 165-169 (1987) に発表.

^{**} 東京生化学研究所