

Title	A simplified synthesis of 32-oxygenated lanosterol derivatives
Sub Title	
Author	園田, よし子(Sonoda, Yoshiko) 田上, 淑恵(Tanoue, Yoshie) 山口, 真起子(Yamaguchi, Makiko) 佐藤, 良博(Sato, Yoshihiro)
Publisher	共立薬科大学
Publication year	1987
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.32 (1987.) ,p.62- 62
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000032-0062

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

A Simplified Synthesis of 32-Oxygenated Lanosterol Derivatives*

Yoshiko SONODA, Yoshie TANOUE, Makiko YAMAGUCHI and Yoshihiro SATO

園田よし子, 田上淑恵, 山口真起子, 佐藤良博

The biosynthesis of cholesterol from lanosterol requires the removal of the three methyl groups at carbons 4 and 14. The initial step in the removal of these methyl groups has been considered to be the 14-demethylation, which is a complex process, and many aspects of the overall mechanisms remain unclear. A probable intermediate is the lanosterol derivatives with a 14-hydroxymethyl, 14-aldehyde or 15-hydroxy group. With the intention of investigating the effects of the natural precursors on cholesterol biosynthesis from 24,25-dihydrolanosterol, we studied the synthesis of the 32-oxygenated derivatives of 24,25-dihydrolanosterol. This report describes a simplified synthesis of lanost-8-ene-3 β ,32-diol, lanost-7-ene-3 β ,32-diol, 3 β -hydroxylanost-8-en-32-al, and 3 β -hydroxylanost-7-en-32-al.

* 本報告は *Chem. Pharm. Bull.*, **35**, 394—397 (1987) に発表.