Title	Constituents of the seed of malva verticillata. III. characterization of the major pectic peptidoglycan and oligosaccharides
Sub Title	
Author	権田, 良子(Gonda, Ryoko) 友田, 正司(Tomoda, Masashi) 金成, 美枝子(Kanari, Mieko) 米田, 直美(Yoneda, Naomi) 清水, 十和子(Shimizu, Towako) 重城, 啓子(Shigeshiro, Keiko) 矢沢, 知美(Yazawa, Tomomi)
Publisher	共立薬科大学
Publication year	1988
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.33 (1988. ) ,p.138- 138
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000033- 0138

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって 保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

## Constituents of the Seed of *Malva verticillata*. III. Characterization of the Major Pectic Peptidoglycan and Oligosaccharides\*

Ryoko Gonda, Masashi Tomoda, Mieko Kanari, Naomi Yoneda, Towako Shimizu, Keiko Shigeshiro, and Tomomi Yazawa

> 権田良子,友田正司,金成美枝子,米田直美, 清水十和子,重城啓子,矢沢知美

The major pectic peptidoglycan, designated as MVS-V, was isolated by chromatography of hot water extract with DEAE-Sephadex A-25 from the seeds of *Malva verticillata* L. Gel chromatography gave a value of 22000 for the molecular weight.

MVS-V is composed of 42.8% polysaccharide and 57.2% protein moieties. The polysaccharide contained L-arabinose, D-xylose, L-rhamnose, D-galactose, and D-galacturonic acid, and the molar ratio was 6:5:8:3:24. In adition, MVS-V contains 0.7% methoxyl and 1.4% O-acetyl groups. Thus one per six galacturonic acid residues in MVS-V exists as methyl esters.

Neither measurement of NMR spectrum nor methylation analysis of MVS-V gave good results. Thus, the carbohydrate moiety in MVS-V was isolated by treatment with a protease followed by gel chromatography. The polysaccharide fraction obtained was composed of 89.6% carbohydrate moiety having the same sugar composition as that of MVS-V, and 10.4% peptide. This fraction is tentatively designated as MVS-V-CH. Methylation analysis of the carboxyl-reduced derivative of MVS-V-CH, <sup>13</sup>C-NMR spectrum of MVS-V-CH, and periodate oxidation studies of MVS-V suggested that the minimal repeating unit of MVS-V-CH is composed of three terminal  $\alpha$ -L-arabinofuranose, three 1,5-linked  $\alpha$ -L-arabinofuranose, five 1,4-linked  $\beta$ -D-xylopyranose, one terminal  $\beta$ -D-galactopyranose, two 3,4-branched  $\beta$ -Dgalactopyranose, six 1,2-linked  $\alpha$ -L-rhamnopyranose, two 2,4-branched  $\alpha$ -Lrhamnopyranose, and twenty-four 1,4-linked  $\alpha$ -D-galacturonic acid residues. As a product of partial hydrolysis of MVS-V,  $2-O-\alpha-(D-galactopyranosyluronic$ acid) - L-rhamnopyranose was identified.

It can be concluded that the polysaccaride moiety of MVS-V resembles that of pectic substances. It has a main chain consisting of  $\alpha$ -1,4-linked D-galacturonic acid residues which are interspersed with  $\alpha$ -1,2-linked L-rhamnose residues, and a quarter of the rhamnose residues occupy branching points bearing arabinoxylogalactan side chains at position 4.

On the other hand, several oligosaccharides belonging to the "raffinose family" were isolated from the hot water extract of the material seeds. Stachyose, raffinose, sucrose, and in addition, D-fructose and D-glucitol were identified.

<sup>\*</sup> Chem. Pharm. Bull., 36, 2790-2795 (1988) に発表.