Title	Copper (II) chelates of an intermediate species absorbing in the 500-nm region
Sub Title	
Author	松島, 美一(Matsushima, Yoshikazu)
	永田, 佳子(Nagata, Yoshiko)
Publisher	共立薬科大学
Publication year	1988
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of
	Pharmacy). No.33 (1988. ) ,p.142- 142
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000033-0142

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって 保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

## Copper (II) Chelates of an Intermediate Species Absorbing in the 500-nm Region\*

Yoshikazu MATSUSHIMA and Yoshiko NAGATA

松島美一,永田佳子

A key step in the action of almost all pyridoxal enzymes is thought to be the formation of a quinoid structure, in which the  $\alpha$ -carbon of the pyridoxylideneamino acid (aldimine) is deprotonated. Several enzymes have been reported to exhibit an intense absorption band in the 500-nm region of the spectrum on addition of substrate or pseudosubstrate. This absorption has been ascribed to the quinoid intermediate. We previously observed an intermediate species absorbing at 488 nm in Al(III) chelate mediated nonenzymatic reaction between a ketimine from ethyl pyruvate and pyridoxamine and an aldimine from ethyl alaninate and pyridoxal. It was identified as the quinoid intermediate. Other trivalent ions of the aluminum subgroup (Ga, In) and of most lanthanide produced similar intermediates. It has been believed that divalent ions such as Cu(II) do not form the species in the reaction.

Our recent study showed that an intermediate absorbing in the 500-nm region was formed in a nonenzymatic transamination of pyridoxal and amino acid ester catalyzed by the Cu(II) chelate of dipyridyl or tripyridyl or related ligands. The absorption was ascribed to the ternary complex of Cu(II)-quinoid-ligand. The quinoid species may be stabilized in a six-coordinated ternary Cu(II) complex.

<sup>\*</sup> 本報告は "Biochemistry of Vitamin B<sub>6</sub>" ed. by T.Korpela and P.Christen, Birkhäuser, Basel, 1987, pp. 367-370 に発表.