

Title	Chromatographic separation and identification of human urine components
Sub Title	
Author	永田, 佳子(Nagata, Yoshiko) 松島, 美一(Matsushima, Yoshikazu) 鵜飼, 晴美(Ugai, Harumi) 菊池, 真弓(Kikuchi, Mayumi) 高井, 信治(Takai, Nobuharu)
Publisher	共立薬科大学
Publication year	1988
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.33 (1988. ) ,p.145- 145
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	<a href="https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000033-0145">https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000033-0145</a>

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

## Chromatographic Separation and Identification of Human Urine Components\*

Yoshiko NAGATA, Yoshikazu MATSUSHIMA, Harumi UGAI,  
Mayumi KIKUCHI, and Nobuharu TAKAI\*\*

永田佳子, 松島美一, 鷗飼晴美, 菊池真弓, 高井信治\*\*

We reported previously the analysis of ultraviolet (UV) -absorbing compounds in human urine by means of high-performance liquid chromatography (HPLC) using the column of macroreticular anion exchange resin (Diaion CDR-10) maintained at 60 °C. The method is expected to find its place among routine analytical methods of human urine applicable to diagnostic purpose. In the study, the chromatographic peaks were assigned by the comparison of the retention times with the authentic samples. For further studies, the separation and identification of the urine components are required.

The peak fractions were collected with a large scale column (250 × 28 mm I.D. stainless-steel) of Diaion CDR-10. The sample was eluted with a linear gradient of 0.006 M to 6.0 M ammonium acetate buffer (pH, 4.40) and the eluates corresponding to several peaks were collected. The reversed phase HPLC of fractions showed that they were almost free of UV-absorbing contaminants. A compound purified from a fraction was subjected to spectral studies. Its UV, infrared and mass spectral data were almost superimposable to those of the authentic sample of uric acid. Thus the compound was identified as uric acid.

The fractionation through HPLC with a large scale column, the purification and the identification by physicochemical means were successful for human urine components.

---

\* 本報告は *Chromatographia*, 24, 317-318 (1987) に発表.

\*\* 東京大学生産技術研究所