

| | |
|------------------|---|
| Title | High-performance liquid chromatographic analysis of ginsenosides in panax ginseng extracts using glass-ODS column |
| Sub Title | |
| Author | 金沢, 秀子(Kanazawa, Hideko) 永田, 佳子(Nagata, Yoshiko) 松島, 美一(Matsushima, Yoshikazu) 友田, 正司(Tomodara, Masashi) 高井, 信治(Takai, Nobuharu) |
| Publisher | 共立薬科大学 |
| Publication year | 1988 |
| Jtitle | 共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.33 (1988.) ,p.146- 146 |
| JaLC DOI | |
| Abstract | |
| Notes | 抄録 |
| Genre | Technical Report |
| URL | https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000033-0146 |

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

High-Performance Liquid Chromatographic Analysis of Ginsenosides in *Panax Ginseng* Extracts Using Glass-ODS Column*

Hideko KANAZAWA, Yoshiko NAGATA, Yoshikazu MATSUSHIMA,
Masashi TOMODA, and Nobuharu TAKAI**

金沢秀子, 永田佳子, 松島美一, 友田正司, 高井信治**

A high-performance liquid chromatographic (HPLC) method using a column of octadecylsilyl porous glass (glass-ODS) achieved rapid and excellent separation. As a part of our interest in HPLC analysis of biological and medical materials, we applied the method for analysis of ginsenosides.

A mixture of ginsenosides was analyzed by the chromatography with 203 nm detection. Ginsenosides R_f, R_g₂, R_b₁, R_c, R_b₂, and R_d were separated with acetonitrile-water (27.5 : 72.5) as the mobile phase. Also a well-resolved chromatogram of ginsenosides R_o, R_g₁ and R_e was obtained with acetonitrile-water (16.5 : 83.5). The whole separation was achieved in 12 min with a flow-rate of 1 ml/min. Calibration curves of ginsenosides R_b₁, R_c, R_b₂, R_d, R_g₁ and R_e were linear up to 5 μg.

Ginseng was extracted with methanol and the extract was filtered, dried and refrigerated. Separation of the ginsenoside peaks of the extract was quite satisfactory. The ginsenoside fractions were taken and subjected to the HPLC analysis. Each fraction gave a single chromatographic peak. The fractions were also analyzed by means of the thin layer chromatography. They gave single spots and showed the same R_f values as the standard samples of the assigned peaks.

The present method using glass-ODS column has an advantage in rapid separation of ginsenosides and will find its place among other analytical methods which are in routine use.

* 本報告は *Chromatographia*, 24, 517-519 (1987) に発表.

** 東京大学生産技術研究所