Title	A comment on the evaluation of equilibrium constants for α -tocopherol interactions with fatty acids by absorbance in the ultraviolet region
Sub Title	
Author	菅田, 節朗(Sugata, Setsuro) 浦野, 四郎(Urano, Shiro) 松島, 美一(Matsushima, Yoshikazu) 松尾, 光芳(Matsuo, Mitsuyoshi)
Publisher	共立薬科大学
Publication year	1988
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.33 (1988.),p.147- 147
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000033-0147

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

**A comment on the evaluation of equilibrium constants for
 α -tocopherol interactions with fatty acids by
absorbance in the ultraviolet region***

Setsuro SUGATA, Shiro URANO,** Yoshikazu MATSUSHIMA,
and Mitsuyoshi MATSUO**

菅田節朗, 浦野四郎**, 松島美一, 松尾光芳**

Vitamin E, especially α -tocopherol, has been considered to act as a biological antioxidant and/or a membrane stabilizer in the lipid core of biomembranes. It is of great importance, therefore, to study the interaction of α -tocopherol and lipids for the verification of membrane stabilization by vitamin E.

Erin and coworkers measured the spectral changes of α -tocopherol in ethanol solutions by the addition of a fatty acid (Erin, A.N., Spirin, M.M., Tabidze, L.V. and Kagan, V.E. (1984) *Biochim. Biophys. Acta* 774, 96—102). They calculated equilibrium constants for α -tocopherol interactions with the fatty acid on the basis of the findings that the absorbance of α -tocopherol in the 200-nm region decreased with an increase in the concentration of the fatty acid.

In the studies on vitamin E-related compounds, we applied their method and obtained the almost same results. During the course of the investigation, the question was raised whether their method provided the reliable constants in a solvent used. Thus, we reexamined their method.

There seems to be no theoretical basis for their assumption that the decrease in the absorption maxima at the 200-nm region indicates an α -tocopherol interaction with a fatty acid.

A quantitative treatment of spectral data is valid only in the case of solutions which do not show a considerable absorbance. The solvent and fatty acids used in their work have considerable absorbance in the 200-nm region of the spectra.

On these basis, it is concluded that the ultraviolet method is inadequate, because it is affected by the absorption in the region of the solvent, ethanol, and fatty acids which they used.

* 本報告は *Biochim. Biophys. Acta*, 962, 385—386 (1988) に発表.

** 老人総合研究所