

Title	In vitro effects of oxygenated lanosterol derivatives on cholesterol biosynthesis from 24, 25-dihydrolanosterol
Sub Title	
Author	園田, よし子(Sonoda, Yoshiko) 関川, 善夫(Sekigawa, Yoshio) 佐藤, 良博(Sato, Yoshihiro)
Publisher	共立薬科大学
Publication year	1988
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.33 (1988.) ,p.159- 159
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000033-0159

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

In Vitro* Effects of Oxygenated Lanosterol Derivatives on Cholesterol Biosynthesis from 24,25-Dihydrolanosterol

Yoshiko SONODA, Yoshio SEKIGAWA, and Yoshihiro SATO

園田よし子, 関川善夫, 佐藤良博

The effects of oxygenated lanosterol derivatives ($5\ \mu\text{M}$) including 3 α -oxygenated lanosterol derivatives on cholesterol biosynthesis from [24,25- ^3H]-24,25-dihydrolanosterol ($18\ \mu\text{M}$) were tested in $10000\times g$ supernatant (S-10) fraction of rat liver homogenate. Among the derivatives, 7-oxolanost-8-en-3 β -ol (7-oxo-DHL), 3 β -acetylanost-8-en-7-one (7-oxo-DHL-3-OAc), and 7-oxolanosta-5,8,11-trien-3 β -ol were highly active in depressing cholesterol biosynthesis from 24,25-dihydrolanosterol. The inhibitory activities of these derivatives on cholesterol synthesis are discussed on the basis of the position and stereochemistry of the oxygen functional groups on the sterol nucleus. The effect of aphidicolin on cholesterol synthesis was also compared with that of 7-oxo-DHL.

* 本報告は *Chem. Pharm. Bull.*, **36**, 966–973 (1988) に発表.