

Title	Metabolism of 32-hydroxylated 24, 25-dihydrolanosterols by partially purified cytochrome P-450 _{14DM} from rat liver microsome
Sub Title	
Author	関川, 善夫(Sekigawa, Yoshio) 園田, よし子(Sonoda, Yoshiko) 佐藤, 良博(Sato, Yoshihiro)
Publisher	共立薬科大学
Publication year	1988
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.33 (1988.) ,p.160- 160
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000033-0160

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Metabolism of 32-Hydroxylated 24,25-Dihydrolanosterols by Partially Purified Cytochrome P-450_{14DM} from Rat Liver Microsome*

Yoshio SEKIGAWA, Yoshiko SONODA, and Yoshihiro SATO

関川善夫, 園田よし子, 佐藤良博

Metabolism of 32-hydroxylated 24,25-dihydrolanosterols, including the intermediate of lanosterol and 24,25-dihydrolanosterol (DHL) demethylation, were studied in a reconstituted system consisting of rat liver partially purified cytochrome P-450, which catalyzes lanosterol 14-demethylation (P-450_{14DM}), and NADPH-cytochrome P-450 reductase. The reconstituted system converted lanost-8-ene-3 β ,32-diol (Δ^8 -32-OH) to 4,4-dimethyl-5 α -cholesta-8,14-dien-3 β -ol, the 14-dehydroxymethylated product, in the same way as DHL. Lanost-7-ene-3 β ,32-diol and lanost-6-ene-3 β ,32-diol, the isomers of Δ^8 -32-OH, were not converted to the corresponding 14-dehydroxymethylated products. The apparent K_m value of P-450_{14DM} for Δ^8 -32-OH was about 1/6 of that for DHL. The metabolism of DHL was inhibited by 7-oxo-24,25-dihydrolanosterol (7-oxo-DHL), which is a potent inhibitor of cholesterol biosynthesis from lanosterol or DHL. However, the metabolism of Δ^8 -32-OH was less inhibited by 7-oxo-DHL than that of DHL.

* 本報告は *Chem. Pharm. Bull.*, **36**, 3049-3054 (1988) に発表.