

Title	Carbon 13 nuclear magnetic resonance study of meso-hexestrol and its derivatives
Sub Title	
Author	小田, 泰子(Oda, Taiko) 村井, 知子(Murai, Tomoko) 佐藤, 良博(Sato, Yoshihiro)
Publisher	共立薬科大学
Publication year	1988
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.33 (1988.) ,p.161- 161
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000033-0161

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Carbon-13 Nuclear Magnetic Resonance Study of *meso*-Hexestrol and Its Derivatives*

Taiko ODA, Tomoko MURAI, and Yoshihiro SATO

小田泰子, 村井知子, 佐藤良博

The carbon-13 nuclear magnetic resonance chemical shift assignments for *meso*-hexestrol, made on the basis of two-dimensional ^{13}C - ^1H chemical shift correlation, long-range selective ^1H decoupling experiment, and a reported two-dimensional Fourier-transform experiment for long-range proton-carbon-13 spin coupling constants, are reported. For measurement of carbon-proton coupling constants of *meso*-hexestrol derivatives, the coupling information was detected by using a gated decoupling facility which permitted retention of the nuclear Overhauser enhancement and a long-range selective ^1H decoupling experiment. The results showed that the aromatic carbon resonances are influenced by the structure (no double bond, or one or two double bond(s) of the hexane framework in the central portion.

* 本報告は *Chem. Pharm. Bull.*, 36, 1534-1539 (1988) に発表.