

Title	Studies on the biosynthesis of bialaphos (SF-1293) 8 purification and characterization of 2-phosphinomethylmalic acid synthase from streptomyces hygroscopicus SF-1293
Sub Title	
Author	下遠野, 久美子(Shimotono, Kumiko) 瀬戸, 治男(Seto, Haruo) 大岳, 望(Otake, Noboru) 今井, 敏(Imai, Satoshi) 村上, 健(Murakami, Takeshi)
Publisher	共立薬科大学
Publication year	1988
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.33 (1988. ) ,p.172- 172
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	<a href="https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000033-0172">https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000033-0172</a>

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

**Studies on the Biosynthesis of Bialaphos (SF-1293) & Purification  
and Characterization of 2-Phosphinomethylmalic acid  
Synthase from *Streptomyces hygroscopicus* SF-1293\***

Kumiko W. SHIMOTOHNO, Haruo SETO\*\*, Noboru ÔTAKE\*\*,  
Satoshi IMAI\*\*\* and Takeshi MURAKAMI\*\*\*\*

下遠野久美子, 瀬戸治男\*\*, 大岳 望\*\*  
今井 敏\*\*\*, 村上 健\*\*\*\*

2-Phosphinomethylmalic acid (PMM) synthase catalyzes the condensation of phosphinopyruvic acid (PPA), an analog of oxalacetic acid, and acetyl-CoA to form PMM. The enzyme was purified approximately 700-fold from a cell-free extract of *Streptomyces hygroscopicus* SF-1293, a bialaphos producing organism, to an electrophoretically homogeneous state. The purified PMM synthase has a subunit molecular weight of 48,000 by SDS-polyacrylamide gel electrophoresis and a native molecular weight of 90,000 ~ 98,000 by gel filtration. PMM synthase was relatively unstable, showed maximum activity at pH 8.0 and 30°C, and was inhibited strongly by *p*-chloromercuribenzoate, iodoacetamide and EDTA. Enzyme activity suppressed by EDTA was completely restored by adding Co<sup>++</sup> or Mn<sup>++</sup> and partially restored by addition of Ca<sup>++</sup>, Fe<sup>++</sup> or Mg<sup>++</sup>.

The specific substrates of this enzyme are PPA or oxalacetic acid in addition to acetyl-CoA. The enzyme does not catalyze the liberation of CoA from acetyl-CoA in the presence of  $\alpha$ -keto acids, such as pyruvate,  $\alpha$ -ketoglutarate, deamino- $\alpha$ -keto-demethylphosphinothricin or phosphonopyruvate. The condensation reaction did not take place when propionyl-CoA or butyryl-CoA was used as a substrate in place of acetyl-CoA. The Km values of the enzyme were 0.05 mM for acetyl-CoA, 0.39 mM for PPA and 0.13 mM for oxalacetate. PMM synthase is very similar to (*R*)-citrate synthase of *Clostridium* in the inhibition pattern by sulfhydryl compounds, its metal ion requirement and stereospecificity; unlike (*R*)-citrate synthase PMM synthase was not inhibited by oxygen.

\* 本報告は *J. Antibiotics*, 41, 1057-1065 (1988) に発表.

\*\* 東京大学応微研

\*\*\* 明治製菓薬開研

\*\*\*\* 明治製菓薬品研