慶應義塾大学学術情報リポジトリ

Keio Associated Repository of Academic resouces

| Title | Determination of air luminescence spectra for alpha-emitters with liquid scintillation spectrometers |
|------------------|--|
| Sub Title | |
| Author | 村瀬, 裕子(Murase, Yuko) 本間, 義夫(Honma, Yoshio) 滝上, 誠(Takiue, Makoto) 油井, 多丸(Aburai, Tamaru) |
| Publisher | 共立薬科大学 |
| Publication year | 1988 |
| Jtitle | 共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.33 (1988.) ,p.173- 173 |
| JaLC DOI | |
| Abstract | |
| Notes | 沙 録 |
| Genre | Technical Report |
| URL | https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000033-0173 |

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって 保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Determination of Air Luminescence Spectra for Alpha-Emitters with Liquid Scintillation Spectrometers

Yuko Murase, Yoshio Homma, Makoto Takiue**, and Tamaru Aburai***

村瀬裕子,本間義夫,滝上 誠**,油井多丸***

Air luminescence spectra have been measured for 210 Po, 238 U and 241 Am with two commercially available liquid scintillation spectrometers. Since the measured endpoint of the air luminescence spectra is approx. 18 keV, they can be determined by liquid scintillation counting in the 3 H detection region. The average counting efficiencies measured with both counting systems were found to be 33.3% and 33.7%, respectively. The air luminescence effect on liquid scintillation counting of gaseous α -emitters is mentioned.

^{*} 本報告は Appl. Radiat. Isot. (in press)

^{**} 慈恵会医科大学

^{***} 日本原子力研究所ラジオアイソトープ・原子炉研修所.