Title	Anticomplementary and hypoglycemic activity of okra and hibiscus mucilages
Sub Title	
Author	友田, 正司(Tomodota, Masashi) 清水, 訓子(Shimizu, Noriko) 権田, 良子(Gonda, Ryoko) 金成, 美枝子(Kanari, Mieko) 山田, 陽城(Yamada, Haruki) ヒキノ, ヒロシ(Hikino, Hiroshi)
Publisher	共立薬科大学
Publication year	1989
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.34 (1989.),p.58- 58
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000034-0058

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Anticomplementary and Hypoglycemic Activity of Okra and Hibiscus Mucilages*

Masashi TOMODA, Noriko SHIMIZU, Ryōko GONDA, Mieko KANARI,
Haruki YAMADA** and Hiroshi HIKINO***

友田正司, 清水訓子, 権田良子, 金成美枝子, 山田陽城**,
ヒキノヒロシ***

The present paper describes the anticomplementary activity and hypoglycemic activity of the six mucilages obtained from *Abelmoschus esculentus* (i.e. *Hibiscus esculentus* L.; okra), *Hibiscus moscheutos* and *Hibiscus syriacus*. Okra-mucilages F and R were isolated from the immature fruit and the root of *Abelmoschus esculentus*, respectively. Hibiscus-mucilages Mo and ML were isolated from the root and the leaf of *Hibiscus moscheutos*, and Hibiscus-mucilages SL and SF from the leaf and the flower bud of *Hibiscus syriacus*, respectively.

Hibiscus-mucilages ML and SF showed remarkable anticomplementary activities. Okra-mucilage R and Hibiscus-mucilage SL also had potent activities, which were almost at the same level as that of the positive control, AR-arabinogalactan from the root of *Angelica acutiloba*. The remarkable activities of these mucilages may result from the highly branched structure. The activities of Okra-mucilage F and Hibiscus-mucilage Mo were lower than those of the former four mucilages.

Most of Okra-mucilages F and R and of Hibiscus-mucilages Mo, ML, SL and SF exhibited significant hypoglycemic activity, though the effect of Hibiscus-mucilage ML was dose dependent. These mucilages which showed extensive hypoglycemic activity have a backbone chain consisting of almost only the repeating unit L-rhamno-D-galacturonan. Okra-mucilage R especially showed a remarkable activity. It possesses the trisaccharide repeating unit, $(1 \rightarrow 4)\text{-}[\beta\text{-D-Glc}pA\text{-}(1 \rightarrow 3)]\text{-}\alpha\text{-D-GalpA}\text{-}(1 \rightarrow 2)\text{-}\alpha\text{-L-Rhap}$, as the main part of the backbone. In addition, it has side chains composed mainly of $\beta\text{-1,4-linked D-galactopyranose}$ residues, and $\alpha\text{-1,2-linked L-rhamnopyranosyl}$ residues in part of the backbone. The D-glucuronic acid side-chain groups at O-3 of the D-galacturonic acid residues in the backbone are not essential for the hypoglycemic effect, as Okra-mucilage F showed a significantly activity.

* 本報告は *Carbohydr. Res.*, **190**, 323—328 (1989) に発表。

** 北里研究所東洋医学総研

*** 東北大学薬学部