

| | |
|------------------|---|
| Title | Anticomplementary and hypoglycemic activity of okra and hibiscus mucilages |
| Sub Title | |
| Author | 友田, 正司(Tomododa, Masashi) 清水, 訓子(Shimizu, Noriko) 権田, 良子(Gonda, Ryoko) 金成, 美枝子(Kanari, Mieko) 山田, 陽城(Yamada, Haruki) ヒキノ, ヒロシ(Hikino, Hiroshi) |
| Publisher | 共立薬科大学 |
| Publication year | 1989 |
| Jtitle | 共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.34 (1989.),p.58- 58 |
| JaLC DOI | |
| Abstract | |
| Notes | 抄録 |
| Genre | Technical Report |
| URL | https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000034-0058 |

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Anticomplementary and Hypoglycemic Activity of Okra and Hibiscus Mucilages*

Masashi TOMODA, Noriko SHIMIZU, Ryōko GONDA, Mieko KANARI,
Haruki YAMADA** and Hiroshi HIKINO***

友田正司, 清水訓子, 権田良子, 金成美枝子, 山田陽城**,
ヒキノヒロシ***

The present paper describes the anticomplementary activity and hypoglycemic activity of the six mucilages obtained from *Abelmoschus esculentus* (i.e. *Hibiscus esculentus* L.; okra), *Hibiscus moscheutos* and *Hibiscus syriacus*. Okra-mucilages F and R were isolated from the immature fruit and the root of *Abelmoschus esculentus*, respectively. Hibiscus-mucilages Mo and ML were isolated from the root and the leaf of *Hibiscus moscheutos*, and Hibiscus-mucilages SL and SF from the leaf and the flower bud of *Hibiscus syriacus*, respectively.

Hibiscus-mucilages ML and SF showed remarkable anticomplementary activities. Okra-mucilage R and Hibiscus-mucilage SL also had potent activities, which were almost at the same level as that of the positive control, AR-arabinogalactan from the root of *Angelica acutiloba*. The remarkable activities of these mucilages may result from the highly branched structure. The activities of Okra-mucilage F and Hibiscus-mucilage Mo were lower than those of the former four mucilages.

Most of Okra-mucilages F and R and of Hibiscus-mucilages Mo, ML, SL and SF exhibited significant hypoglycemic activity, though the effect of Hibiscus-mucilage ML was dose dependent. These mucilages which showed extensive hypoglycemic activity have a backbone chain consisting of almost only the repeating unit L-rhamno-D-galacturonan. Okra-mucilage R especially showed a remarkable activity. It possesses the trisaccharide repeating unit, $(1 \rightarrow 4) - [\beta\text{-D-Glc}pA - (1 \rightarrow 3)] - \alpha\text{-D-Gal}pA - (1 \rightarrow 2) - \alpha\text{-L-Rhap}$, as the main part of the backbone. In addition, it has side chains composed mainly of β -1,4-linked D-galactopyranose residues, and α -1,2-linked L-rhamnopyranosyl residues in part of the backbone. The D-glucuronic acid side-chain groups at O-3 of the D-galacturonic acid residues in the backbone are not essential for the hypoglycemic effect, as Okra-mucilage F showed a significantly activity.

* 本報告は *Carbohydr. Res.*, **190**, 323—328 (1989) に発表。

** 北里研究所東洋医学総研

*** 東北大学薬学部