Title	A reticuloendothelial system-activating glycan from the seeds of malva verticillata
Sub Title	
Author	友田, 正司(Tomododa, Masashi) 金成, 美枝子(Kanari, Mieko) 権田, 良子(Gonda, Ryoko) 清水, 訓子(Shimizu, Noriko)
Publisher	共立薬科大学
Publication year	1989
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.34 (1989.),p.59- 59
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000034-0059

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

A Reticuloendothelial System-Activating Glycan from the Seeds of *Malva verticillata**

Masashi TOMODA, Mieko KANARI, Ryōko GONDA and Noriko SHIMIZU

友田正司, 金成美枝子, 権田良子, 清水訓子

From the hot water extract of the seeds of *Malva verticillata*, a polysaccharide, designated as MVS-III A, has been isolated by fractionation on DEAE-Sephadex A-25 (carbonate) followed by gel chromatography on Sephacryl S-500 and affinity chromatography on Con A-Sepharose columns. The polysaccharide gave a single band on PAGE and gave a single peak on gel chromatography.

MVS-III A is composed of L-arabinose : D-xylose : D-galactose : D-galacturonic acid in the molar ratio of 16 : 1 : 8 : 3, and it contains a peptide moiety (1.7%). Its value of molecular mass was estimated to be about 8.5×10^6 .

The results of methylation analysis of the original polysaccharide and the carboxyl-reduced derivative and ^{13}C -NMR spectrum suggested that the minimal unit of the polysaccharide is composed of four terminal α -L-arabinofuranose, ten α -1,5-linked L-arabinofuranose, two α -1,3-linked L-arabinopyranose, one β -1,4-linked D-xylopyranose, four β -1,3-linked D-galactopyranose, four 3,6-branched β -D-galactopyranose and α -1,4-linked D-galacturonan residues. The periodate oxidation studies also supported these conclusions.

The effect of MVS-III A on a RES was demonstrated by the *in vivo* carbon clearance test. When administered i.p. (50 mg/kg), the phagocytic index of MVS-III A was 0.4492 ± 0.1048 . Thus the value was quite remarkably increased, suggesting powerful activation of RES by i.p. injection of MVS-III A.

The structural characterization of MVS-III A is both its high arabinose content and the presence of α -1,3-linked L-arabinopyranose residues. Most of the RES-activating polysaccharides from Oriental crude drugs contain arabino-3,6-branched galactan units in common as their major parts, though they have different types of backbone chain.

* 本報告は *Phytochemistry* 28, 2609—2611 (1989) に発表。