

Title	Separation and purification of uridine diphosphate-glucuronosyltransferases by a chromatofocusing on a high-performance liquid chromatograph
Sub Title	
Author	高梨, 浩典(Takanashi, Hironori) 本間, 浩(Honma, Hiroshi) 松井, 道夫(Matsui, Michio)
Publisher	共立薬科大学
Publication year	1989
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.34 (1989.) ,p.65- 65
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000034-0065

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Separation and Purification of Uridine Diphosphate-Glucuronosyltransferases by a Chromatofocusing on a High-Performance Liquid Chromatograph*

Hironori TAKANASHI, Hiroshi HOMMA and Michio MATSUI

高梨浩典, 本間 浩, 松井道夫

A rapid method for the separation and purification of uridine diphosphate-glucuronosyltransferases (GT) was developed with the use of chromatofocusing on high-performance liquid chromatograph. GT isoenzymes solubilized from hepatic microsomes of Wistar rats were separated on a Mono P column, a pre-packed column for chromatofocusing. Using 4-nitrophenol, testosterone and androsterone as substrates, four fractions with different GT activities were separated in a pH gradient from 9.5 to 7.0. Two isoenzymes, testosterone GT and androsterone GT were purified to apparent homogeneity. They were eluted at pH 8.9 and 8.0 and had subunit molecular weight values of 50000 and 52000, respectively. Approximately 10 mg of solubilized microsomal proteins were applied and the elution was completed within 2 hr. Addition of N-nitrodiethylamine, an *in vitro* activator of GT activity, enhanced the GT activity toward 4-nitrophenol in the three fractions. This chromatographic analysis confirmed the absence of androsterone GT isoenzyme in LA Wistar rats, a mutant strain in terms of androsterone glucuronidation.

* 本報告は *Chem. Pharm. Bull.*, 37, 1583—1586 (1989) に発表.