Title	Continuous separation of glucosides in dual-flow extraction system
Sub Title	
Author	西沢, 秀幸(Nishizawa, Hideyuki) 渡辺, 右佳子(Watanabe, Yukako) 沖村, 早希子(Okimura, Sakiko) 阿部, 芳廣(Abe, Yoshihiro)
Publisher	共立薬科大学
Publication year	1989
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.34 (1989.) ,p.67- 67
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000034- 0067

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって 保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Continuous Separation of Glucosides in Dual-Flow Extraction System*

Hideyuki NISHIZAWA, Yukako WATANABE, Sakiko Okimura and Yoshihiro Abe

西沢秀幸,渡辺右佳子,沖村早希子,阿部芳廣

A new separation system in dual-flow countercurrent extraction has been established. As a result, a two-component mixture could be separated with the improved apparatus, liquid particle extractor, whose working principle is the same as that in liquid particle chromatography (LPC). Liquid particle chromatography has been used as a means to determine the distribution constants of components in a sample, which were necessary for setting the operating condition of the extractor. The optimum condition for the best separation of two compounds (distribution constant of D_1 and D_2) was deduced theoretically. Thus, the flow rate of the aqueous phase (H) could be determined by a calculation using the flow rates of the sample feed (A, fed as an organic solution) and the organic phase (S) by the relation : $H=(S+A/2)\sqrt{D_1\times D_2}$. With the extractor, a cpouinuous and complete separation of phenyl β -D-glucopyranoside (D=1.17) and esculin (D=1.58) in a 100 mg-scale (100 mg each in 20 ml of water saturated 1-butanol) could be executed in 14 h. A precise separation of a mixture with a separation iactor (D_1/D_2) of 1.23 was also accomplished.

^{*} 本報告は Anal. Sci., 5 (3), 345—350 (1989) に発表.