Title	Identification of cholesta-7, 24-dien-3 β -ol and desmosterol in hamster cauda epididymal spermatozoa
Sub Title	
Author	粟野, みのり(Awano, Minori) 川口, 昭彦(Kawaguchi, Akihiko) 森崎, 益雄(Morisaki, Masuo) 毛利, 秀雄(Mori, Hideo)
Publisher	共立薬科大学
Publication year	1989
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.34 (1989.), p.84- 84
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000034-0084

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

**Identification of Cholesta-7,24-dien-3 β -ol and Desmosterol
in Hamster Cauda Epididymal Spermatozoa***

Minori AWANO**, Akihiko KAWAGUCHI**, Masuo MORISAKI
and Hideo MOHRI**

栗野みのり**, 川口昭彦**, 森崎益雄, 毛利秀雄**

The sterol composition of hamster cauda epididymal spermatozoa was remarkably different from that of several other mammalian spermatozoa. Desmosterol and cholesta-7,24-dien-3 β -ol account for as much as 90% of the total sterols. Cholesterol and desmosterol are the major components of mouse cauda epididymal spermatozoa, and rabbit, boar and bull ejaculated spermatozoa. Cholesta-7,24-dien-3 β -ol was not detected. Furthermore, cholesterol was the main sterol in hamster caput epididymal spermatozoa, while only a trace amount of desmosterol was detected and cholesta-7,24-dien-3 β -ol was hardly detected at all. The sterol content of cauda and caput epididymal spermatozoa was $0.17 \pm 0.05 \mu\text{mol}/10^8$ spermatozoa. During maturation, the desmosterol and cholesta-7,24-dien-3 β -ol levels increase and the cholesterol level decreases. Cholesta-7,4-dien-3 β -ol appears as a sterol in mature spermatozoa and seems to be a characteristic sterol of hamster cauda epididymal spermatozoa.

---

* 本報告は *Lipids* 24, 662—664 (1989) に発表.

** 東京大学教養学部