Title	Surface potential of mixed micelles composed of prostaglandin B ₂ and heptaethylene glycol dodecyl ether by the fluorescence method
Sub Title	
Author	横山, 祥子(Yokoyama, Shoko) 金子, 明子(Kaneko, Akiko) 藤江, 忠雄(Fujie, Tadao)
Publisher	共立薬科大学
Publication year	1989
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.34 (1989.),p.90- 90
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000034-0090

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Surface Potential of Mixed Micelles Composed of Prostaglandin B₂ and Heptaethylene Glycol Dodecyl Ether by the Fluorescence Method*

Shoko YOKOYAMA, Akiko KANEKO, and Tadao FUJIE

横山祥子, 金子明子, 藤江忠雄

The surface state of negatively charged mixed micelles prepared by addition of prostaglandin B₂ (PGB₂) to nonionic heptaethylene glycol dodecyl ether (HED) has been investigated and the surface potential ($\Delta\psi$) of the micelles has been estimated by measuring the fluorescence intensity of ammonium 8-anilino-1-naphthalenesulfonate (ANS) as a fluorescent probe. Since the binding constant of ANS, k , is dependent on the surface potential of the micelle, the value of $\Delta\psi$ was determined according to the equation $\Delta\psi = -59.16 \log(k_0/k)$ at 25°C, where k_0 is the k of HED micelle. The value of $|\Delta\psi|$ increases with increasing mole fraction of PGB₂ (X_{PGB_2}) and decreases with increasing ionic strength (J). These changes are due to the effects of the negative charge of PGB₂ and of ionic atmosphere of Na⁺, respectively. Surface charge density (σ) was, therefore, determined from $\Delta\psi$ according to the Gouy-Chapman theory, with a result that σ is nearly independent of J and proportional to X_{PGB_2} . These results have led to a conclusion that the state of the diffuse layer surrounding the Stern layer on the micelle surface is represented well by the Gouy-Chapman theory as well as a previous result for the system of sodium dodecyl sulfate (SDS) and HED. The value of $|\sigma|$ for PGB₂ micelle is less than that for SDS micelle, which suggests that PGB₂ forms a loosely packed micelle.

* 本報告は *Bull. Chem. Soc. Jpn.*, **61** (10), 3451—3454 (1988) に発表.