

Title	Characterization of polysaccharides having activity on the reticuloendothelial system from the rhizome of curcuma longa
Sub Title	
Author	権田, 良子(Gonda, Ryoko) 友田, 正司(Tomododa, Masashi) 清水, 訓子(Shimizu, Noriko) 金成, 美枝子(Kanari, Mieko)
Publisher	共立薬科大学
Publication year	1990
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.35 (1990.) ,p.34- 34
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000035-0034

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

**Characterization of Polysaccharides Having Activity
on the Reticuloendothelial System from
the Rhizome of *Curcuma longa****

Ryōko GONDA, Masashi TOMODA, Noriko SHIMIZU and Mieko KANARI

権田良子, 友田正司, 清水訓子, 金成美枝子

The crude polysaccharide fraction was isolated from the rhizome of *Curcuma longa* by hot water extraction followed by precipitation with ethanol. It was fractionated by DEAE-Sephadex A-25 (carbonate) column chromatography, and the fraction having high activity was subjected to affinity chromatography on Con A-Sepharose and gel chromatography on Sephacryl S-300. Three polysaccharides, named ukonan A, ukonan B and ukonan C were obtained. They were homogeneous on electrophoresis and gel chromatography, and showed remarkable reticuloendothelial system (RES)-potentiating activity in a carbon clearance test even in a minute dose (5 mg per kg).

Among them, ukonan B is the major one in terms of the activity. It contains L-arabinose, D-xylose, D-galactose, D-glucose, L-rhamnose and D-galacturonic acid in the molar ratio of 12:4:12:1:2:4 in addition to 6.6% peptide moiety. The value of 3.3×10^4 was estimated for its molecular mass.

Methylation analysis and ^{13}C -NMR studies indicated that the minimal unit of ukonan B is composed of twenty-four terminal α -L-arabinofuranose, ten α -1,3-linked L-arabinopyranose, two α -1,5-linked L-arabinofuranose, two β -1,4-linked D-xylose, ten β -3,4-branched D-xylose, four terminal β -D-galactose, ten β -1,3-linked D-galactose, four β -1,6-linked D-galactose, eighteen β -3,6-branched D-galactose, two terminal α -D-glucose, one α -1,4-linked D-glucose, four α -1,2-linked L-rhamnose, two α -2,4-branched L-rhamnose and twelve α -1,4-linked D-galacturonic acid residues. Thus, ukonan B may be basically a kind of α -L-arabino- β -3,6-branched D-galactan.

* 本報告は *Chem. Pharm. Bull.*, 38, 482-486 (1990) に発表.