Title	Structure and anticomplementary activity of an acidic polysaccharide from the leaves of malva sylvestris var. mauritiana
Sub Title	
Author	権田, 良子(Gonda, Ryoko) 友田, 正司(Tomododa, Masashi) 清水, 訓子(Shimizu, Noriko) 山田, 陽城(Yamada, Haruki)
Publisher	共立薬科大学
Publication year	1990
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.35 (1990.),p.38- 38
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000035-0038

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

**Structure and Anticomplementary Activity of an Acidic
Polysaccharide from the Leaves of *Malva
sylvestris* var. *mauritiana****

Ryōko GONDA, Masashi TOMODA, Noriko SHIMIZU and Haruki YAMADA**

権田良子, 友田正司, 清水訓子, 山田陽城**

A main polysaccharide, designated as MSL-P, was isolated from the leaves of *Malva sylvestris* var. *mauritiana* by extraction with water followed by fractionation with DEAE-Sephadex A-25 (carbonate) and purification by gel chromatography with Sephacryl S-300 and Sephadex G-25. It was homogeneous by electrophoresis and gel chromatography, which gave a value of 1.1×10^4 as molecular mass.

It is composed of L-rhamnose, D-galactose, D-galacturonic acid and D-glucuronic acid in the molar ratio of 22:6:22:11, and it contains 7.7% peptide moiety.

The structure of the polysaccharide component was elucidated by methylation analysis, partial hydrolysis and ^{13}C -NMR studies. The backbone of the polysaccharide is composed of alternating α -1,4-linked D-galacturonic acid and α -1,2-linked L-rhamnosyl residues. Approximately 50% of the 1,4-linked D-galacturonic acid residues are also substituted at position 3 with terminal, non-reducing β -D-glucuronic acid groups. Some of the 1,2-linked L-rhamnosyl residues are also substituted at position 4 with oligosaccharides containing β -1,4-linked D-galactosyl residues.

MSL-P had potent anti-complementary activity, which was nearly equal to that of the positive control, AR-4, from the root of *Angelica acutiloba*.

* 本報告は *Carbohydr. Res.*, **198**, 323—329 (1990) に発表.

** 北里研究所東医総研