

Title	Characterization of two polysaccharides having activity on the reticuloendothelial system from the root of glycyrrhiza uralensis
Sub Title	
Author	友田, 正司(Tomododa, Masashi) 清水, 訓子(Shimizu, Noriko) 金成, 美枝子(Kanari, Mieko) 権田, 良子(Gonda, Ryoko) 新井, 澄子(Arai, Sumiko) 奥田, 有子(Okuda, Yuko)
Publisher	共立薬科大学
Publication year	1990
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.35 (1990.) ,p.39- 39
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000035-0039

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

**Characterization of Two Polysaccharides Having Activity
on the Reticuloendothelial System from the
Root of *Glycyrrhiza uralensis****

Masashi TOMODA, Noriko SHIMIZU, Mieko KANARI, Ryōko GONDA,
Sumiko ARAI and Yūko OKUDA

友田正司, 清水訓子, 金成美枝子, 権田良子, 新井澄子, 奥田有子

Two polysaccharides called glycyrrhizans UA and UB, were isolated from the root of *Glycyrrhiza uralensis* by hot water extraction followed by treatment with cetyltrimethylammonium bromide, gel chromatography on Sephacryl S-300 and affinity chromatography on Con A-Sepharose. Each polysaccharide gave a single band on PAGE, and gave a single peak on gel chromatography. They showed significant reticuloendothelial system-potentiating activity in a carbon clearance test.

Glycyrrhizan UA is composed of L-arabinose: D-galactose: L-rhamnose: D-galacturonic acid in the molar ratio of 20:14:1:3, and glycyrrhizan UB is composed of L-arabinose: D-galactose: D-glucose: L-rhamnose: D-galacturonic acid in the molar ratio of 12:10:1:10:20, in addition to small amounts of O-acetyl groups (2.9% in UA and 2.5% in UB) and peptide moiety. About 10% (UA) and 35% (UB) of the D-galacturonic acid residues exist as the methyl esters. Their molecular masses were estimated to be 6.9×10^4 and 1.07×10^4 .

Chemical and spectroscopic studies indicated that the minimal unit of glycyrrhizan UA is composed of twenty-two terminal α -L-arabinofuranose, twenty-six α -1,5-linked L-arabinofuranose, four terminal α -L-arabinopyranose, eight α -1,3-linked L-arabinopyranose, one terminal β -D-galactopyranose, eight β -1,3-linked D-galactopyranose, eight β -1,4-linked D-galactopyranose, three β -1,6-linked D-galactopyranose, twenty-two β -3,6-branched D-galactopyranose, one α -1,2-linked L-rhamnopyranose, one α -2,4-branched L-rhamnopyranose, six α -1,4-linked D-galactopyranosyluronic acid and three α -2,4-branched D-galactopyranosyluronic acid residues. The minimal unit of glycyrrhizan UB is composed of the same four, four arabinofuranose, one, three arabinopyranose, three, two, one, one, two galactopyranose, six, four rhamnose, eighteen and two galacturonic acid residues as those in glycyrrhizan UA, in addition to one β -2,4-branched D-galactopyranose and one terminal α -D-glucopyranose residues.

* 本報告は *Chem. Pharm. Bull.*, 38, 1667—1671 (1990) に発表.