

Title	Ca ²⁺ -insensitive contraction in glycerinated skeletal muscle fiber treated with additional troponin T
Sub Title	
Author	中山, 雪麿(Nakayama, Yukimaro) 青木, 裕美(Aoki, Hiromi) 渡辺, 和子(Watanabe, Kazuko) 山口, 正弘(Yamaguchi, Masahiro)
Publisher	共立薬科大学
Publication year	1990
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.35 (1990.) ,p.45- 45
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000035-0045

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Ca²⁺-insensitive Contraction in Glycerinated Skeletal Muscle Fiber Treated with Additional Troponin T*

Yukimaro NAKAYAMA, Hiromi AOKI, Kazuko WATANABE**
and Masahiro YAMAGUCHI***

中山雪麿, 青木裕美, 渡辺和子**, 山口正弘***

An investigation was carried out on the effect of troponin T (TnT) protein on the Ca²⁺-sensitivity of fiber contraction when it was added to glycerinated muscle fiber. The TnT-treated fiber was observed to lose its Ca²⁺-sensitivity, so that it was seen to contract even in a Ca²⁺ free buffer solution. However, the treated fibers were not significantly different from the normally glycerinated ones in terms of tension magnitude.

On the other hand, the fiber did not lose its Ca²⁺-sensitivity upon treatment in a similar manner with troponin I (TnI) or troponin C (TnC) protein instead of TnT; the tension magnitude was seen to decrease only with TnI treatment.

SDS-PAGE patterns of the TnT-treated fibers were different from those of glycerinated ones at the TnI and TnC bands, which were fainter in the TnT-treated fibers than in the normal-ones. DACM-labeled fibers also decreased in the DACM-labeled TnI and TnC proteins on gel as the result of the treatment with TnT protein. Further, those proteins were completely removed from fibers into extra medium.

These findings suggest that either the additional TnT protein removes TnI and TnC subunits from the Tn complex combined with thin filaments in the fibers, or that it exchanges the total Tn complex on the thin filaments.

* 本報告は *Jpn. J. Physiol.* 40, 403-410 (1990) に発表.

** 日本大学・薬学部・生物学

*** 順天堂大学・体育学部・栄養生化学