

| | |
|------------------|---|
| Title | Radioactivity in deuterated compounds |
| Sub Title | |
| Author | 本間, 義夫(Honma, Yoshio) 村瀬, 裕子(Murase, Yuko) |
| Publisher | 共立薬科大学 |
| Publication year | 1990 |
| Jtitle | 共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.35 (1990.) ,p.49- 49 |
| JaLC DOI | |
| Abstract | |
| Notes | 抄録 |
| Genre | Technical Report |
| URL | https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000035-0049 |

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Radioactivity in Deuterated Compounds*

Yoshio HOMMA and Yuko MURASE

本間義夫, 村瀬裕子

Commercially available D₂O and deuterated toluene have been measured with a liquid scintillation spectrometer and were found to contain 2.71—6.15 Bq of tritium in 1 ml of D₂O, and 1.10 Bq of tritium in 1 ml of deuterated toluene.

* 本報告は *J. Radioanal. Nucl. Chem. Letters* **146**(3), 205—210 (1990) に発表.