

Title	Temperature dependence of the pulse-height distributions in aliphatic liquid scintillators
Sub Title	
Author	村瀬, 裕子(Murase, Yuko) 本間, 義夫(Honma, Yoshio)
Publisher	共立薬科大学
Publication year	1990
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.35 (1990. ) ,p.50- 50
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	<a href="https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000035-0050">https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000035-0050</a>

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

## Temperature Dependence of the Pulse-Height Distributions in Aliphatic Liquid Scintillators\*

Yuko MURASE and Yoshio HOMMA

村瀬裕子, 本間義夫

The pulse-height distributions for  $^{131\text{m}}\text{Xe}$  in PPO solutions of aliphatic hydrocarbons such as n-pentane, n-hexane, n-heptane, 2,4-dimethylpentane, 2,3,4-trimethylpentane, 1-pentene, 1-hexene and 1-heptene are investigated as a function of temperature. The pulse-height distributions are found to be shifted toward higher pulse-height with decreasing temperature. The count rates of  $^{131\text{m}}\text{Xe}$  remain unchanged with decreasing temperature. The mechanism of the effect is also discussed.

---

\* 本報告は *J. Radioanal. Nucl. Chem. Letters* **146**(4), 215—222 (1990) に発表.