

Title	Pharmacological differentiation of presynaptic M <sub>1</sub> muscarinic receptors modulating acetylcholine release from postsynaptic muscarinic receptors in guinea-pig ileum
Sub Title	
Author	川島, 紘一郎(Kawashima, Koichiro) 藤本, 和子(Fujimoto, Kazuko) 鈴木, 岳之(Suzuki, Takeshi) 大畑, 尚代(Ohata, Hisayo)
Publisher	共立薬科大学
Publication year	1990
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.35 (1990. ) ,p.61- 61
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	<a href="https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000035-0061">https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000035-0061</a>

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

**Pharmacological Differentiation of Presynaptic M<sub>1</sub> Muscarinic Receptors Modulating Acetylcholine Release from Postsynaptic Muscarinic Receptors in Guinea-Pig Ileum\***

Koichiro KAWASHIMA, Kazuko FUJIMOTO, Takeshi SUZUKI and Hisayo OOHATA

川島紘一郎, 藤本和子, 鈴木岳之, 大畑尚代

1. Effects of three muscarinic antagonists on electrically evoked ACh release and contractile response were investigated in longitudinal muscle strips of guinea-pig ileum suspended in an organ-bath and superfused with Krebs solution. ACh release was determined by a specific radioimmunoassay.

2. Telenzepine, a selective M<sub>1</sub> muscarinic antagonist, increased the ACh release at a concentration of 100-fold less than that inhibiting the contractile response (10 vs 1000 nM).

3. AF-DX 116, a cardioselective M<sub>2</sub> muscarinic antagonist, inhibited the contractile response at 10 μM, but did not affect the ACh release at this concentration.

4. (–)N-Methylscopolamine (NMS) did not affect the ACh release, but inhibited the contractile response at all concentrations tested (1–1000 nM), indicating (–)NMS can be used as an ileal specific postsynaptic muscarinic antagonist.

5. These data demonstrate that presynaptic muscarinic receptors modulating ACh release are distinct from postsynaptic ones involved in the contractile response and can be classified as M<sub>1</sub> subtype.

---

\* 本報告は *Gen. Pharmac* Vol. 21, No. 1, pp. 17–21, 1990 に発表.