Title	Hemicholinium-3-resistant choline uptake system linked to acetylcholine synthesis in the rat hippocampus
Sub Title	
Author	鈴木, 岳之(Suzuki, Takeshi) 鹿島, 裕子(Kashima, Yuko) 川島, 紘一郎(Kawashima, Koichiro)
Publisher	共立薬科大学
Publication year	1990
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.35 (1990.),p.63- 63
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000035-0063

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Hemicholinium-3-resistant Choline Uptake System Linked to Acetylcholine Synthesis in the Rat Hippocampus*

Takeshi SUZUKI, Yuko KASHIMA and Koichiro KAWASHIMA

鈴木岳之, 鹿島裕子, 川島紘一郎

Extracellular choline-dependency of acetylcholine (ACh) synthesis was examined in rat hippocampal slices. In the presence of hemicholinium-3 (HC-3, 5 μ M), extracellular choline-dependent ACh synthesis appeared to consist of two different components. The first component (saturated at up to 50 μ M choline) was dependent on the HC-3-resistant choline uptake system having a low capacity for choline in comparison with the high-affinity choline uptake system (HACU). The second component, observed in the presence of more than 50 μ M choline, was considered due to competitive inhibition of HACU by HC-3. HACU-dependent ACh synthesis seemed to be saturated in the presence of up to 2 μ M choline. These results indicate that both the HACU and HC-3-resistant choline uptake systems are linked to ACh synthesis in the hippocampus

* 本報告は *Neuroscience Letters*, 105 (1989) 211—214 に発表.