

Title	Simultaneous determination of ginsenosides and saikosaponins by HPLC
Sub Title	
Author	金沢, 秀子(Kanazawa, Hideko) 永田, 佳子(Nagata, Yoshiko) 松島, 美一(Matsushima, Yoshikazu) 友田, 正司(Tomoda, Masashi) 高井, 信治(Takai, Nobuharu)
Publisher	共立薬科大学
Publication year	1990
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.35 (1990. ) ,p.68- 68
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	<a href="https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000035-0068">https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000035-0068</a>

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

## Simultaneous Determination of Ginsenosides and Saikosaponins by HPLC\*

Hideko KANAZAWA, Yoshiko NAGATA, Yoshikazu MATSUSHIMA,  
Masashi TOMODA and Nobuharu TAKAI\*\*

金沢秀子, 永田佳子, 松島美一, 友田正司, 高井信治\*\*

We prepared octadecylsilyl porous glass (MPG-ODS) as the packing for reversed phase high-performance liquid chromatography (HPLC) and achieved rapid and excellent separation and quantitative determination of ginsenosides with the column. Ginsenosides are saponins of ginseng, the roots of *Panax ginseng*, which has long been used in traditional oriental medicine.

In the medicine, several crude drugs are generally prescribed in a single formula. Among the important crude drugs often prescribed with ginseng are bupleurum root, the roots of *Bupleurum falcatum*, and glycyrrhiza. The pharmacological studies on these crude drugs have centered on the saponins. Saikosaponins and glycyrrhizin were isolated from bupleurum root and glycyrrhiza, respectively. Since the crude drugs are used in the same pharmaceutical preparations, simultaneous determination of the saponins were often required.

A mixture of ginsenosides and saikosaponins were analyzed by HPLC using the column of MPG-ODS with detection at 203 nm. A well-resolved chromatogram of ginsenoside Rb<sub>1</sub>, Rc, Rb<sub>2</sub>, Rd and saikosaponin a, b<sub>2</sub> and c was obtained with acetonitrile-water (25.5: 74.5) as the mobile phase. The whole separation was achieved in 20 min with a flow-rate of 1.5 ml/min. Calibration curves of ginsenoside Rb<sub>1</sub>, Rc, Rb<sub>2</sub>, Rd and saikosaponin a, c were linear up to 5 µg. A pharmaceutical preparation of commercial origin, "Shosaiko-to", was analyzed by the procedure. It was concluded that the rapid and accurate simultaneous determinations of the saponins is possible by the described method.

---

\* 本報告は *J. Chromatogr.*, **507**, 327—332 (1990) に発表.

\*\* 東京大学生産技術研究所.