

| | |
|------------------|---|
| Title | Preparative high-performance liquid chromatography on chemically modified porous glass. isolation of saponins from ginseng |
| Sub Title | |
| Author | 金沢, 秀子(Kanazawa, Hideko) 永田, 佳子(Nagata, Yoshiko) 松島, 美一(Matsushima, Yoshikazu) 友田, 正司(Tomoda, Masashi) 高井, 信治(Takai, Nobuharu) |
| Publisher | 共立薬科大学 |
| Publication year | 1990 |
| Jtitle | 共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.35 (1990.) ,p.69- 69 |
| JaLC DOI | |
| Abstract | |
| Notes | 抄録 |
| Genre | Technical Report |
| URL | https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000035-0069 |

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

**Preparative High-Performance Liquid Chromatography
on Chemically Modified Porous Glass.
Isolation of Saponins from Ginseng*.**

Hideko KANAZAWA, Yoshiko NAGATA, Yoshikazu MATSUSHIMA,
Masashi TOMODA and Nobuharu TAKAI**

金沢秀子, 永田佳子, 松島美一, 友田正司, 高井信治**

Preparative high-performance liquid chromatography on octadecylsilyl porous glass (pore size, 550Å) with acetonitrile-water as the mobile phase was applied for the isolation of saponins from *Panax ginseng*. An automatic fraction collection system was used. In a single run, several milligrams of pure ginsenosides were obtained from 10 grams of roots of *Panax ginseng*. The excellent separation may be partly due to the optimal pore size of 550 Å and the narrow distribution range of the size. The composition of organic solvent in the mobile phase was much smaller in HPLC with MPG-ODS than with silica-ODS columns. The savings in organic solvents are advantageous for both economical and environmental reasons. The method was simple, rapid and convenient and should be applicable to isolation of other saponins of crude drugs.

* 本報告は *Chem. Pharm. Bull.*, 33, 1630—1632 (1990) に発表.

** 東京大学生産技術研究所