

Title	Association of octadecanoic acid in 1, 2-dichloroethane
Sub Title	
Author	横山, 祥子(Yokoyama, Shoko) 藤江, 忠雄(Fujie, Tadao)
Publisher	共立薬科大学
Publication year	1990
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.35 (1990. ) ,p.75- 75
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	<a href="https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000035-0075">https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000035-0075</a>

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

## Association of Octadecanoic Acid in 1,2-Dichloroethane\*

Shoko Yokoyama and Tadao Fujie

横山祥子, 藤江忠雄

The association of octadecanoic acid (C18) in 1,2-dichloroethane at 40°C has been investigated by measuring the fluorescence spectra and intensities of ammonium 8-anilino-1-naphthalenesulfonate as a probe.

As a result, it has been found that C18 begins to associate at a concentration of  $1.0\text{--}1.2 \times 10^{-2}$  mol dm<sup>-3</sup> and that the aggregation number is small and constant in the concentration region above about  $4 \times 10^{-2}$  mol dm<sup>-3</sup>.

The mechanism for the formation of fatty acid (FA)-thiamine disulfide (TDS) complexes, (FA)<sub>6</sub>(TDS), was also discussed.

---

\* 本報告は *Chem. Pharm. Bull.*, 38 (8), 2249—2252 (1990) に発表.